

10/23/00
JC949 U.S. PTO

10-25-00

Express Mail Label No. EL497498851US
Date of Deposit: October 23, 2000
File No.: 00-53

Box Seq

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
APPLICATION FEE TRANSMITTAL

Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

Sir:

Transmitted herewith for filing under 37 CFR §1.53(b) is the

- ☒ patent application ☐ continuation patent application
☐ divisional patent application ☐ continuation-in-part patent application

of Applicant: Debra G. Gilbertson

Title: METHOD FOR TREATING FIBROSIS

- ☒ 45 pages of specification ☒ 8 sheets of drawings
☒ 14 pages of sequence listing
☐ An assignment of the invention to _____
☒ 2 sheets of ☐ signed ☒ unsigned Declaration and Power of Attorney
☒ **ASCII Computer Disk Sequence pursuant to 37 C.F.R. 1.821(f). It is believed that the content of the paper sequence listing and the computer readable sequence listing are the same.**

CALCULATION OF APPLICATION FEE

Claim Type	No. Filed	Less	Extra	Extra Rate	Fee
Total	18	-20	0	\$18.00	\$000.00
Independent	3	-3	0	\$80.00	\$000.00
Basic Fee					\$710.00
Multiple Dependency Fee If Applicable (\$270.00)					\$000.00
Total Filing Fee					\$710.00

- ☐ Priority of application Serial No. _____ filed on _____ in _____ is claimed under 35 U.S.C. 119. A certified copy thereof is submitted herewith.


- ☒ The benefit of applications Serial Nos. 60/161,653 filed on October 21, 1999; Serial No. 60/165,255 filed on November 12, 1999; and Serial No. 60/222,223 filed on August 1, 2000; all in the U.S. Patent and Trademark Office, is claimed under 35 U.S.C. 120 or 119(e) 1.

Please charge ZymoGenetics, Inc., Deposit Account No. 26-0290 as follows:

- ☒ Filing fee, estimated to be \$710.00
☐ Assignment recording fee
☒ Any additional fees associated with this paper or during the pendency of this application.
☐ The issue fee set in 37 C.F.R. 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

A copy of this sheet is enclosed.

Respectfully submitted,



Gary E. Parker
Registration No. 31,648

JC914 U.S. PTO
09/695121
10/23/00

09695121 102300

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

EXPRESS MAIL CERTIFICATE

Box Patent Application
Commissioner for Patents
Washington, DC 20231

Re: U.S. Patent Application for
METHOD FOR TREATING FIBROSIS

Applicant: Debra G. Gilbertson

Sir:


Express Mail Label No. EL497498851US

Date of Deposit October 23, 2000

I hereby certify that the following attached paper(s) or fee

1. Return Post card
2. Application Fee Transmittal (in duplicate)
3. Patent Application (45 pages)
4. Figures 1-2 (8 sheets of drawings)
5. Unexecuted Declaration and Power of Attorney
6. Sequence Listing (14 pages)
7. ASCII Computer Disk Sequence pursuant to 37 CFR 1.821(e).

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 C.F.R. 1.10 on the date indicated above, addressed to the Commissioner for Patents, Washington, DC 20231.


Amy Toman

ZymoGenetics, Inc
1201 Eastlake Avenue East
Seattle, WA 98102
(206) 442-6600



09/695121

Variable	Mean	SD	Min	Max
Age	34.2	10.5	18	65
Gender	0.5	0.5	0	1
Marital Status	0.6	0.5	0	1
Education	12.5	1.5	9	16
Income	3500	1500	1000	8000
Health	0.8	0.2	0	1
Stress	4.5	1.5	1	7
Depression	0.3	0.4	0	1
Life Satisfaction	5.5	1.5	1	9
Resilience	0.7	0.3	0	1
Optimism	0.6	0.4	0	1
Gratitude	0.5	0.5	0	1
Forgiveness	0.4	0.5	0	1
Empathy	0.6	0.4	0	1
Compassion	0.5	0.5	0	1
Kindness	0.4	0.5	0	1
Generosity	0.3	0.5	0	1
Patience	0.5	0.5	0	1
Humility	0.4	0.5	0	1
Modesty	0.3	0.5	0	1
Self-control	0.6	0.4	0	1
Discipline	0.5	0.5	0	1
Perseverance	0.4	0.5	0	1
Endurance	0.3	0.5	0	1
Stamina	0.4	0.5	0	1
Strength	0.3	0.5	0	1
Power	0.2	0.5	0	1
Influence	0.3	0.5	0	1
Authority	0.2	0.5	0	1
Leadership	0.3	0.5	0	1
Management	0.2	0.5	0	1
Organization	0.3	0.5	0	1
Planning	0.2	0.5	0	1
Decision-making	0.3	0.5	0	1
Problem-solving	0.2	0.5	0	1
Communication	0.3	0.5	0	1
Interpersonal Skills	0.2	0.5	0	1
Teamwork	0.3	0.5	0	1
Collaboration	0.2	0.5	0	1
Cooperation	0.3	0.5	0	1
Partnership	0.2	0.5	0	1
Relationship	0.3	0.5	0	1
Connection	0.2	0.5	0	1
Network	0.3	0.5	0	1
Community	0.2	0.5	0	1
Society	0.3	0.5	0	1
Culture	0.2	0.5	0	1
Tradition	0.3	0.5	0	1
Customs	0.2	0.5	0	1
Religion	0.3	0.5	0	1
Spirituality	0.2	0.5	0	1
Philosophy	0.3	0.5	0	1
Worldview	0.2	0.5	0	1
Beliefs	0.3	0.5	0	1
Values	0.2	0.5	0	1
Principles	0.3	0.5	0	1
Standards	0.2	0.5	0	1
Criteria	0.3	0.5	0	1
Guidelines	0.2	0.5	0	1
Rules	0.3	0.5	0	1
Laws	0.2	0.5	0	1
Regulations	0.3	0.5	0	1
Procedures	0.2	0.5	0	1
Protocols	0.3	0.5	0	1
Methods	0.2	0.5	0	1
Techniques	0.3	0.5	0	1
Skills	0.2	0.5	0	1
Abilities	0.3	0.5	0	1
Talents	0.2	0.5	0	1
Gifts	0.3	0.5	0	1
Strengths	0.2	0.5	0	1
Weaknesses	0.3	0.5	0	1
Assets	0.2	0.5	0	1
Liabilities	0.3	0.5	0	1
Resources	0.2	0.5	0	1
Capabilities	0.3	0.5	0	1
Potential	0.2	0.5	0	1
Opportunities	0.3	0.5	0	1
Challenges	0.2	0.5	0	1
Risks	0.3	0.5	0	1
Uncertainties	0.2	0.5	0	1
Unknowns	0.3	0.5	0	1
Variables	0.2	0.5	0	1
Factors	0.3	0.5	0	1
Elements	0.2	0.5	0	1
Components	0.3	0.5	0	1

OF

FOR

METHOD FOR TREATING FIBROSIS

Description

METHOD OF TREATING FIBROSIS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C. 119(e) from U.S. provisional applications Serial Nos. 60/161,653, filed October 21, 1999; 60/165,255, filed November 12, 1999; and 60/222,223, filed August 1, 2000.

BACKGROUND OF THE INVENTION

Fibrosis is the abnormal accumulation of fibrous tissue that can occur as a part of the wound-healing process in damaged tissue. Such tissue damage may result from physical injury, inflammation, infection, exposure to toxins, and other causes. Examples of fibrosis include dermal scar formation, keloids, liver fibrosis, lung fibrosis (e.g., silicosis, asbestosis), kidney fibrosis (including diabetic nephropathy), and glomerulosclerosis.

Liver (hepatic) fibrosis, for example, occurs as a part of the wound-healing response to chronic liver injury. Fibrosis occurs as a complication of haemochromatosis, Wilson's disease, alcoholism, schistosomiasis, viral hepatitis, bile duct obstruction, exposure to toxins, and metabolic disorders. This formation of scar tissue is believed to represent an attempt by the body to encapsulate the injured tissue. Liver fibrosis is characterized by the accumulation of extracellular matrix that can be distinguished qualitatively from that in normal liver. Left unchecked, hepatic fibrosis progresses to cirrhosis (defined by the presence of encapsulated nodules), liver failure, and death.

In recent years there have been significant advances in the understanding of the cellular and biochemical mechanisms underlying liver fibrosis (reviewed by Li and Friedman, *J. Gastroenterol. Hepatol.* 14:618-633, 1999). Stellate cells are believed to be a major source of extracellular matrix in the liver. Stellate cells respond to a variety of cytokines present in the liver, some of which they also produce (Friedman, *Seminars in Liver Disease* 19:129-140, 1999).

As summarized by Li and Friedman (*ibid.*), actual and proposed
35 therapeutic strategies for liver fibrosis include removal of the underlying cause (e.g.,
toxin or infectious agent), suppression of inflammation (using, e.g., corticosteroids, IL-
1 receptor antagonists, or other agents), down-regulation of stellate cell activation

(using, e.g., gamma interferon or antioxidants), promotion of matrix degradation, or promotion of stellate cell apoptosis. Despite recent progress, many of these strategies are still in the experimental stage, and existing therapies are aimed at suppressing inflammation rather than addressing the underlying biochemical processes. Thus, there
5 remains a need in the art for materials and methods for treating fibrosis, including liver fibrosis.

DESCRIPTION OF THE INVENTION

The present invention provides materials and methods for reducing cell
10 proliferation or extracellular matrix production, treating fibrosis, and reducing stellate cell activation in a mammal.

Within one aspect of the invention there is provided a method of reducing cell proliferation or extracellular matrix production in a mammal comprising administering to the mammal a composition comprising a zveg3 antagonist in
15 combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides, in an amount sufficient to reduce cell proliferation or extracellular matrix production. Within certain embodiments of the invention, proliferation of
20 mesangial, endothelial, smooth muscle, fibroblast, osteoblast, osteoclast, stellate, or interstitial cells is reduced. Within other embodiments, the mammal is suffering from a fibroproliferative disorder of the liver, kidney, or bone.

Within another aspect of the present invention there is provided a method of treating fibrosis in a mammal comprising administering to the mammal a
25 composition comprising a therapeutically effective amount of a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides. Within certain embodiment of the invention the fibrosis is liver
30 fibrosis or kidney fibrosis.

Within a third aspect of the invention there is provided a method of reducing stellate cell activation in a mammal comprising administering to the mammal a composition comprising a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group
35 consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides, in an amount sufficient to reduce stellate cell activation. Within one embodiment, the stellate cells are liver stellate cells.

0069541.102300

Within a fourth aspect of the invention there are provided pharmaceutical compositions for use within the above methods. In general, the compositions comprise a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides.

These and other aspects of the invention will become evident upon reference to the following detailed description of the invention and the attached drawings. In the drawings:

Fig. 1 is a Hopp/Woods hydrophilicity profile of the amino acid sequence shown in SEQ ID NO:2. The profile is based on a sliding six-residue window. Buried G, S, and T residues and exposed H, Y, and W residues were ignored. These residues are indicated in the figure by lower case letters.

Fig. 2 is an alignment of human (SEQ ID NO:2) and mouse (SEQ ID NO:4) amino acid sequences.

The term "antagonist" is used herein to denote a compound that reduces a biological activity of another compound. Within the present invention, a "zveg3 antagonist" is a compound that reduces the receptor-mediated biological activity (e.g., mitogenic activity) of zveg3 on a target cell. Antagonists may exert their action by competing with zveg3 for binding sites on a cell-surface receptor, by binding to zveg3 and preventing it from binding to a cell-surface receptor, by otherwise interfering with receptor function, by reducing production of zveg3, or by other means.

"Extracellular matrix" (ECM) is a complex mixture of macromolecules that accumulates within tissues in close apposition to cell surfaces. ECM contains secreted macromolecules such as collagens I, III and IV, fibronectin, laminins, and various proteoglycans. These macromolecules can be organized to provide cohesion to the tissue and can contribute to its structural and mechanical properties. ECM can act as a depository for, and release site of, potent secreted growth factors, and is known to influence growth, survival and differentiation of the cells it surrounds. Pathologic ECM accumulation, if unchecked, can restrict access of nutrients, growth factors and other physiologically important molecules to cells and can lead to the creation of areas of low live cell density. Over time, this accumulation can result in the inability of a tissue to perform its specific metabolic and structural roles, and may ultimately lead to overt cell and tissue death.

An "inhibitory polynucleotide" is a DNA or RNA molecule that reduces or prevents expression (transcription or translation) of a second (target) polynucleotide. Inhibitory polynucleotides include antisense polynucleotides, ribozymes, and external

guide sequences. The term "inhibitory polynucleotide" further includes DNA and RNA molecules that encode the actual inhibitory species, such as DNA molecules that encode ribozymes.

The terms "treat" and "treatment" are used broadly to denote therapeutic and prophylactic interventions that favorably alter a pathological state. Treatments include procedures that moderate or reverse the progression of, reduce the severity of, prevent, or cure a disease.

All references cited herein are incorporated by reference in their entirety.

The present invention provides methods for treating fibrosis in a patient using zvegf3 antagonists. Zvegf3 is a protein that is structurally related to platelet-derived growth factor (PDGF) and the vascular endothelial growth factors (VEGF). This protein has also been designated "VEGF-R" (WIPO Publication WO 99/37671) and, more recently, "PDGF-C" (WO 00/18212). Zvegf3/PDGF-C is a multi-domain protein with significant homology to the PDGF/VEGF family of growth factors. Representative amino acid sequences of human and mouse zvegf3 are shown in SEQ ID NO:2 and SEQ ID NO:4, respectively. DNAs encoding these polypeptides are shown in SEQ ID NOS:1 and 3, respectively.

The term "zvegf3 protein" is used herein to denote proteins comprising the growth factor domain of a zvegf3 polypeptide (e.g., residues 235-345 of human zvegf3 (SEQ ID NO:2) or mouse zvegf3 (SEQ ID NO:4)), wherein said protein is mitogenic for cells expressing cell-surface PDGF α -receptor subunit. Zvegf3 has been found to bind to the $\alpha\alpha$ and $\alpha\beta$ isoforms of PDGF receptor. Zvegf3 proteins include homodimers and heterodimers as disclosed below. Using methods known in the art, zvegf3 proteins can be prepared in a variety of forms, including glycosylated or non-glycosylated, pegylated or non-pegylated, with or without an initial methionine residues, and as fusion proteins as disclosed in more detail below.

Structural predictions based on the zvegf3 sequence and its homology to other growth factors suggests that the polypeptide can form homomultimers or heteromultimers having growth factor activity, i.e., modulating one or more of cell proliferation, migration, differentiation, and metabolism. Experimental evidence confirms that biologically active zvegf3 is a dimeric protein. While not wishing to be bound by theory, the similarity of zvegf3 to other members of the PDGF/VEGF family suggests that zvegf3 may also form heteromultimers with other members of the family, including VEGF, VEGF-B, VEGF-C, VEGF-D, zvegf4 (SEQ ID NO:5), PlGF (Maglione et al., *Proc. Natl. Acad. Sci. USA* 88:9267-9271, 1991), PDGF-A (Murray et

al., U.S. Patent No. 4,899,919; Heldin et al., U.S. Patent No. 5,219,759), or PDGF-B (Chiu et al., *Cell* 37:123-129, 1984; Johnsson et al., *EMBO J.* 3:921-928, 1984).

The zveg3 polypeptide chain comprises a growth factor domain and a CUB domain. The growth factor domain is characterized by an arrangement of cysteine residues and beta strands that is characteristic of the "cystine knot" structure of the PDGF family. The CUB domain shows sequence homology to CUB domains in the neuropilins (Takagi et al., *Neuron* 7:295-307, 1991; Soker et al., *ibid.*), human bone morphogenetic protein-1 (Wozney et al., *Science* 242:1528-1534, 1988), porcine seminal plasma protein and bovine acidic seminal fluid protein (Romero et al., *Nat. Struct. Biol.* 4:783-788, 1997), and *X. laevis* tolloid-like protein (Lin et al., *Dev. Growth Differ.* 39:43-51, 1997).

An alignment of mouse and human zveg3 polypeptide sequences is shown in Fig. 2. Analysis of the amino acid sequence shown in SEQ ID NO:2 indicates that residues 1 to 14 form a secretory peptide. The CUB domain extends from residue 46 to residue 163. A propeptide-like sequence extends from residue 164 to residue 234, and includes two potential cleavage sites at its carboxyl terminus, a dibasic site at residues 231-232 and a target site for furin or a furin-like protease at residues 231-234. The growth factor domain extends from residue 235 to residue 345. Those skilled in the art will recognize that domain boundaries are somewhat imprecise and can be expected to vary by up to ± 5 residues from the specified positions. Potential proteolytic cleavage sites occur at residues 232 and 234. Processing of recombinant zveg3 produced in BHK cells has been found to occur between residues 225 and 226. Signal peptide cleavage is predicted to occur after residue 14 (± 3 residues). This analysis suggests that the zveg3 polypeptide chain may be cleaved to produce a plurality of monomeric species as shown in Table 1. Cleavage after Arg-234 is expected to result in subsequent removal of residues 231-234, with possible conversion of Gly-230 to an amide. Cleavage after Lys-232 is expected to result in subsequent removal of residue 231, again with possible conversion of Gly-230 to an amide. In addition, it may be advantageous to include up to seven residues of the interdomain region at the carboxyl terminus of the CUB domain. The interdomain region can be truncated at its amino terminus by a like amount. See Table 1. Corresponding domains in mouse and other non-human zveg3s can be determined by those of ordinary skill in the art from sequence alignments.

Table 1

<u>Monomer</u>	<u>Residues (SEQ ID NO:2)</u>
Cub domain	15 - 163 46 - 163 15 - 170 46 - 170
CUB domain + interdomain region	15 - 234 46 - 234 15 - 229 amide 15 - 230
Cub domain + interdomain region + growth factor domain	15 - 345 46 - 345
Growth factor domain	235 - 345 226 - 345
Growth factor domain + interdomain region	164 - 345 171 - 345

Zveg3 can thus be prepared in a variety of multimeric forms comprising a zveg3 polypeptide as disclosed above. These zveg3 polypeptides include zveg3₁₅₋₃₄₅, zveg3₄₆₋₃₄₅, zveg3₂₂₆₋₃₄₅, and zveg3₂₃₅₋₃₄₅. Variants and derivatives of these polypeptides can also be prepared as disclosed herein.

Zveg3 proteins can be prepared as fusion proteins comprising amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, an affinity tag, or a targeting polypeptide. For example, a zveg3 protein can be prepared as a fusion with an affinity tag to facilitate purification. In principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag. Affinity tags include, for example, a poly-histidine tract, protein A (Nilsson et al., *EMBO J.* 4:1075, 1985; Nilsson et al., *Methods Enzymol.* 198:3, 1991), glutathione S transferase (Smith and Johnson, *Gene* 67:31, 1988), Glu-Glu affinity tag (Grussenmeyer et al., *Proc. Natl. Acad. Sci. USA* 82:7952-4, 1985), substance P, FlagTM peptide (Hopp et al., *Biotechnology* 6:1204-1210, 1988), streptavidin binding peptide,

maltose binding protein (Guan et al., *Gene* 67:21-30, 1987), cellulose binding protein, thioredoxin, ubiquitin, T7 polymerase, or other antigenic epitope or binding domain. Fusion of zveg3 to, for example, maltose binding protein or glutathione S transferase can be used to improve yield in bacterial expression systems. In these instances the

5 non-zveg3 portion of the fusion protein ordinarily will be removed prior to use. Separation of the zveg3 and non-zveg3 portions of the fusion protein is facilitated by providing a specific cleavage site between the two portions. Such methods are well known in the art. Zveg3 can also be fused to a targetting peptide, such as an antibody (including polyclonal antibodies, monoclonal antibodies, antigen-binding fragments thereof such as F(ab')₂ and Fab fragments, single chain antibodies, and the like) or

10 other peptidic moiety that binds to a target tissue.

Variations can be made in the zveg3 amino acid sequences shown in SEQ ID NO:2 and SEQ ID NO:4 to provide mitogenically inactive, receptor-binding polypeptides that act as zveg3 antagonists. As used herein, the term "mitogenically

15 inactive" means that the protein does not show statistically significant activity in a standard mitogenesis assay as compared to a wild-type zveg3 control. Such variations include amino acid substitutions, deletions, and insertions. While not wishing to be bound by theory, it is believed that residues Arg260-Trp271 of human zveg3 (SEQ ID NO:2) form a loop that define the ability of the protein to bind to PDGF- β receptors,

20 although binding is also permitted to alpha receptors. It is thus predicted that binding to either receptor subunit can be blocked or enhanced by mutations in this region. In addition, residues Leu311-His321 of SEQ ID NO:2 are predicted to form a loop (loop3) that may be mutated to block receptor binding. Peptides that mimic this region of the molecule may act as antagonists.

25 The effects of amino acid sequence changes can be predicted by computer modeling (using, e.g., the Insight II® viewer and homology modeling tools; MSI, San Diego, CA) or determined by analysis of crystal structure (see, e.g., Laphorn et al., *Nature* 369:455, 1994), and can be assessed using art-recognized mutagenesis procedures in combination with activity assays. Representative mutagenesis

30 procedures include, for example, site-directed mutagenesis and alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244, 1081-1085, 1989; Bass et al., *Proc. Natl. Acad. Sci. USA* 88:4498-4502, 1991). Multiple amino acid substitutions can be made and tested using known methods, such as those disclosed by Reidhaar-Olson and Sauer (*Science* 241:53-57, 1988) or Bowie and Sauer (*Proc. Natl. Acad. Sci. USA*

35 86:2152-2156, 1989). Other methods that can be used include phage display (e.g., Lowman et al., *Biochem.* 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204), region-directed mutagenesis

Zveg3 proteins, including full-length polypeptides, fragments, and fusion proteins, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells (including cultured cells of multicellular organisms). Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., eds., *Current Protocols in Molecular Biology*, Green and Wiley and Sons, NY, 1993. See, WO 00/34474. In general, a DNA sequence encoding a zveg3 polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers. See, WO 00/34474.

Zveg3 proteins can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, *cis*-4-hydroxyproline, *trans*-4-hydroxyproline, *N*-methylglycine, *allo*-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipercolic acid, *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-

Zveg3 polypeptides or fragments thereof can also be prepared through chemical synthesis according to methods known in the art, including exclusive solid phase synthesis, partial solid phase methods, fragment condensation or classical solution synthesis. See, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963; Stewart et al., Solid Phase Peptide Synthesis (2nd edition), Pierce Chemical Co., Rockford, IL, 1984; Bayer and Rapp, *Chem. Pept. Prot.* 3:3, 1986; and Atherton et al., *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, Oxford, 1989.

30 Zvegf3 proteins are purified by conventional protein purification methods, typically by a combination of chromatographic techniques. See, in general, Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988; and Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York, 1994. Proteins comprising a polyhistidine affinity tag
35 (typically about 6 histidine residues) are purified by affinity chromatography on a nickel chelate resin. See, for example, Houchuli et al., *Bio/Technol.* 6: 1321-1325, 1988. Furthermore, the growth factor domain itself binds to nickel resin at pH 7.0-8.0

and 25 mM Na phosphate, 0.25 M NaCl. Bound protein can be eluted with a descending pH gradient down to pH 5.0 or an imidazole gradient. Recombinant zveg3 growth factor domain protein can be purified using a combination of chromatography on a strong cation exchanger followed by a tandem column array comprising a strong anion exchanger followed by an immobilized metal affinity column in series. It has also been found that zveg3 binds to various dye matrices (e.g., BLUE1, BLUE 2, ORANGE 1, ORANGE 3, and RED3 from Lexton Scientific, Signal Hill, CA) in PBS at pH 6-8, from which the bound protein can be eluted in 1-2 M NaCl in 20 mM boric acid buffer at pH 8.8. Protein eluted from RED3 may be passed over RED2 (Lexton Scientific) to remove remaining contaminants. Proteins comprising a glu-glu tag can be purified by immunoaffinity chromatography according to conventional procedures. See, for example, Grussenmeyer et al., *ibid.* Maltose binding protein fusions are purified on an amylose column according to methods known in the art.

As disclosed in more detail below, overexpression of zveg3 in the livers of transgenic mice led to marked stellate cell activation and proliferation. At 8 weeks of age there was an accumulation of perisinusoidal extracellular matrix (ECM) that progressed to a perivenular ECM deposition at 22 weeks and possible early stages of cirrhosis characterized by fibrotic banding and hepatic nodule formation at 33 weeks of age. Thus, zveg3 dimers appear to resemble the previously described PDGF isoforms in being potent mitogens of hepatic stellate cells and appearing to play a role in liver fibrosis. These transgenic mice thus provide a model for testing zveg3 antagonists as well as other antifibrotic agents. In view of these and other experiments disclosed herein, it is expected that altered zveg3 expression may initiate or exacerbate a variety of fibrotic conditions. In this context, inhibiting the action of zveg3 using a zveg3 antagonist will limit the progress of such conditions. While not wishing to be bound by theory, it is believed that the pro-fibrotic effects of zveg3 are due at least in part to the induction of TGF- β production.

Zveg3 antagonists include, without limitation, anti-zveg3 antibodies (including neutralizing antibodies), inhibitory polynucleotides (including antisense polynucleotides, ribozymes, and external guide sequences), and other peptidic and non-peptidic agents, including small molecule inhibitors and mitogenically inactive receptor-binding zveg3 polypeptides. Such antagonists can be used to block the mitogenic effects of zveg3 and thereby reduce, inhibit, prevent, or otherwise treat fibrosis, including, without limitation, scar formation, keloids, scleroderma, liver fibrosis, lung fibrosis, kidney fibrosis, pancreatic fibrosis, myelofibrosis, post-surgical fibrotic adhesions, fibroproliferative disorders of the vasculature, fibroproliferative

disorders of the prostate, fibroproliferative disorders of bone, fibromatosis, fibroma, fibrosarcoma, and the like.

Of particular interest is the use of zveg3 or zveg3 antagonists for the treatment or repair of liver damage, including damage due to chronic liver disease, including chronic active hepatitis (including hepatitis C) and many other types of cirrhosis. Widespread, massive necrosis, including destruction of virtually the entire liver, can be caused by, *inter alia*, fulminant viral hepatitis; overdoses of the analgesic acetaminophen; exposure to other drugs and chemicals such as halothane, monoamine oxidase inhibitors, agents employed in the treatment of tuberculosis, phosphorus, carbon tetrachloride, and other industrial chemicals. Conditions associated with ultrastructural lesions that do not necessarily produce obvious liver cell necrosis include Reye's syndrome in children, tetracycline toxicity, and acute fatty liver of pregnancy. Cirrhosis, a diffuse process characterized by fibrosis and a conversion of normal architecture into structurally abnormal nodules, can come about for a variety of reasons including alcohol abuse, post necrotic cirrhosis (usually due to chronic active hepatitis), biliary cirrhosis, pigment cirrhosis, cryptogenic cirrhosis, Wilson's disease, and alpha-1-antitrypsin deficiency. In cases of liver fibrosis it may be beneficial to administer a zveg3 antagonist to suppress the activation of stellate cells, which have been implicated in the production of extracellular matrix in fibrotic liver (Li and Friedman, *ibid.*).

Fibrotic disorders of the kidney include, without limitation, glomerulonephritis (including membranoproliferative, diffuse proliferative, rapidly progressive, and chronic forms), diabetic glomerulosclerosis, focal glomerulosclerosis, diabetic nephropathy, lupus nephritis, tubulointerstitial fibrosis, membranous nephropathy, amyloidosis (which affects the kidney among other tissues), renal arteriosclerosis, and nephrotic syndrome. The glomerulus is a major target of many types of renal injury, including immunologic (e.g., immune-complex- or T-cell-mediated), hemodynamic (systemic or renal hypertension), metabolic (e.g., diabetes), "atherosclerotic" (accumulation of lipids in the glomerulus), infiltrative (e.g., amyloid), and toxic (e.g., snake venom) (Johnson, *Kidney Int.* 45:1769-1782, 1994). The renal structural changes in patients with diabetic nephropathy include hypertrophy of the glomerulus, thickening of the glomerular and tubular membranes (due to accumulated matrix), and increased amounts of matrix in the mesangium and tubulointerstitium (Ziyadeh et al., *Proc. Natl. Acad. Sci. USA* 97:8015-8020, 2000). Glomerular hypertension due to intrarenal hemodynamic changes in diabetes can contribute to the progression of diabetic nephropathy (Ishida et al., *Diabetes* 48:595-602, 1999). Autoimmune nephritis can also lead to altered mesangial cell growth responses (Liu

5

15

20

25

30

5

10

15

35

inoculating a variety of warm-blooded animals such as horses, cows, goats, sheep, dogs, chickens, rabbits, mice, and rats with a zveg3 polypeptide or a fragment thereof.

Immunogenic polypeptides will comprise an epitope-bearing portion of a zveg3 polypeptide (e.g., as shown in SEQ ID NO:2) or receptor. An "epitope" is a region of a protein to which an antibody can bind. See, for example, Geysen et al., *Proc. Natl. Acad. Sci. USA* 81:3998-4002, 1984. Epitopes can be linear or conformational, the latter being composed of discontinuous regions of the protein that form an epitope upon folding of the protein. Linear epitopes are generally at least 6 amino acid residues in length. Relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, Sutcliffe et al., *Science* 219:660-666, 1983. Immunogenic, epitope-bearing polypeptides contain a sequence of at least six, often at least nine, more often from 15 to about 30 contiguous amino acid residues of a zveg3 protein. Polypeptides comprising a larger portion of a zveg3 protein, i.e. from 30 to 50 residues up to the entire sequence are included. It is preferred that the amino acid sequence of the epitope-bearing polypeptide is selected to provide substantial solubility in aqueous solvents, that is the sequence includes relatively hydrophilic residues, and hydrophobic residues are substantially avoided. Such regions include residues 43-48, 96-101, 97-102, 260-265, and 330-335 of SEQ ID NO:2. As noted above, it is generally preferred to use somewhat longer peptides as immunogens, such as a peptide comprising residues 80-104, 299-314, and 299-326 of SEQ ID NO:2. The latter peptide can be prepared with an additional N-terminal Cys residue to facilitate coupling.

The immunogenicity of a polypeptide immunogen may be increased through the use of an adjuvant, such as alum (aluminum hydroxide) or Freund's complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of a zveg3 polypeptide or a portion thereof with an immunoglobulin polypeptide or with maltose binding protein. If the polypeptide portion is "haptten-like", such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), or tetanus toxoid) for immunization.

Alternative techniques for generating or selecting antibodies include *in vitro* exposure of lymphocytes to a polypeptide immunogen, and selection of antibody display libraries in phage or similar vectors (for instance, through use of immobilized or labeled polypeptide). Techniques for creating and screening such random peptide display libraries are known in the art (e.g., Ladner et al., US Patent No. 5,223,409; Ladner et al., US Patent No. 4,946,778; Ladner et al., US Patent No. 5,403,484 and

Ladner et al., US Patent No. 5,571,698), and random peptide display libraries and kits for screening such libraries are available commercially, for instance from Clontech Laboratories (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA), and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ).

- 5 Random peptide display libraries can be screened using the zveg3 sequences disclosed herein to identify proteins which bind to zveg3.

Antibodies are determined to be specifically binding if they bind to their intended target (e.g., zveg3 protein or receptor) with an affinity at least 10-fold greater than the binding affinity to control (e.g., non-zveg3 or non-receptor) polypeptide or protein. In this regard, a "non-zveg3 polypeptide" includes the related molecules VEGF, VEGF-B, VEGF-C, VEGF-D, PlGF, PDGF-A, and PDGF-B, but excludes zveg3 polypeptides from non-human species. Due to the high level of amino acid sequence identity expected between zveg3 orthologs, antibodies specific for human zveg3 may also bind to zveg3 from other species. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, G., *Ann. NY Acad. Sci.* 51: 660-672, 1949). Methods for screening and isolating specific antibodies are well known in the art. See, for example, Paul (ed.), Fundamental Immunology, Raven Press, 1993; Getzoff et al., *Adv. in Immunol.* 43:1-98, 1988; Goding (ed.), Monoclonal Antibodies: Principles and Practice, Academic Press Ltd., 1996; Benjamin et al., *Ann. Rev. Immunol.* 2:67-101, 1984.

Binding affinity can also be determined using a commercially available biosensor instrument (BIAcore™, Pharmacia Biosensor, Piscataway, NJ), wherein protein is immobilized onto the surface of a receptor chip. See, Karlsson, *J. Immunol. Methods* 145:229-240, 1991 and Cunningham and Wells, *J. Mol. Biol.* 234:554-563, 1993. This system allows the determination of on- and off-rates, from which binding affinity can be calculated, and assessment of stoichiometry of binding.

A variety of assays known to those skilled in the art can be utilized to detect antibodies that specifically bind to zveg3 proteins or receptors. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assays, inhibition or competition assays, and sandwich assays.

For therapeutic applications it is generally preferred to use neutralizing antibodies. As used herein, the term "neutralizing antibody" denotes an antibody that inhibits at least 50% of the biological activity of the cognate antigen when the antibody is added at a 1000-fold molar excess. Those of skill in the art will recognize that

greater neutralizing activity is sometimes desirable, and antibodies that provide 50% inhibition at a 100-fold or 10-fold molar excess may be advantageously employed.

Zveg3 antagonists further include antisense polynucleotides, which can be used to inhibit zveg3 gene transcription and thereby inhibit cell activation and/or proliferation *in vivo*. Polynucleotides that are complementary to a segment of a zveg3-encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NO:1) are designed to bind to zveg3-encoding mRNA and to inhibit translation of such mRNA. Antisense polynucleotides can be targetted to specific tissues using a gene therapy approach with specific vectors and/or promoters, such as viral delivery systems as disclosed in more detail below.

Ribozymes can also be used as zveg3 antagonists within the present invention. Ribozymes are RNA molecules that contain a catalytic center and a target RNA binding portion. The term includes RNA enzymes, self-splicing RNAs, self-cleaving RNAs, and nucleic acid molecules that perform these catalytic functions. A ribozyme selectively binds to a target RNA molecule through complementary base pairing, bringing the catalytic center into close proximity with the target sequence. The ribozyme then cleaves the target RNA and is released, after which it is able to bind and cleave additional molecules. A nucleic acid molecule that encodes a ribozyme is termed a "ribozyme gene." Ribozymes can be designed to express endonuclease activity that is directed to a certain target sequence in a mRNA molecule (see, for example, Draper and Macejak, U.S. Patent No. 5,496,698, McSwiggen, U.S. Patent No. 5,525,468, Chowrira and McSwiggen, U.S. Patent No. 5,631,359, and Robertson and Goldberg, U.S. Patent No. 5,225,337). An expression vector can be constructed in which a regulatory element is operably linked to a nucleotide sequence that encodes a ribozyme.

In another approach, expression vectors can be constructed in which a regulatory element directs the production of RNA transcripts capable of promoting RNase P-mediated cleavage of mRNA molecules that encode a zveg3 polypeptide. According to this approach, an external guide sequence can be constructed for directing the endogenous ribozyme, RNase P, to a particular species of intracellular mRNA, which is subsequently cleaved by the cellular ribozyme (see, for example, Altman et al., U.S. Patent No. 5,168,053; Yuan et al., *Science* 263:1269, 1994; Pace et al., WIPO Publication No. WO 96/18733; George et al., WIPO Publication No. WO 96/21731; and Werner et al., WIPO Publication No. WO 97/33991). An external guide sequence generally comprises a ten- to fifteen-nucleotide sequence complementary to zveg3 mRNA, and a 3'-NCCA nucleotide sequence, wherein N is preferably a purine. The external guide sequence transcripts bind to the targeted mRNA species by the formation of base pairs

between the mRNA and the complementary external guide sequences, thus promoting cleavage of mRNA by RNase P at the nucleotide located at the 5'-side of the base-paired region.

5 The growth factor domain of zveg3 has been found to be the active (PDGF receptor-binding) species of the molecule. Proteolytic processing to remove the N-terminal portion of the molecule is required for activation. Thus, inhibitors of this proteolytic activation can also be used as zveg3 antagonists within the present invention.

10 For pharmaceutical use, zveg3 antagonists are formulated for topical or parenteral, particularly intravenous or subcutaneous, delivery according to conventional methods. In general, pharmaceutical formulations will include a zveg3 antagonist in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in water, or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss
15 on vial surfaces, etc. Methods of formulation are well known in the art and are disclosed, for example, in *Remington: The Science and Practice of Pharmacy*, Gennaro, ed., Mack Publishing Co., Easton, PA, 19th ed., 1995. A "therapeutically effective amount" of a composition is that amount that produces a statistically significant effect, such as a statistically significant reduction in disease progression or a
20 statistically significant improvement in organ function. The exact dose will be determined by the clinician according to accepted standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art. The therapeutic formulations will generally be administered over the period required to achieve a beneficial effect,
25 commonly up to several months and, in treatment of chronic conditions, for a year or more. Dosing is daily or intermittently over the period of treatment. Intravenous administration will be by bolus injection or infusion over a typical period of one to several hours. Sustained release formulations can also be employed. For treatment of pulmonary fibrosis, a zveg3 antagonist can be delivered by aerosolization according to
30 methods known in the art. See, for example, Wang et al., US Patent No. 5,011,678; Gonda et al., US Patent No. 5,743,250; and Lloyd et al., US Patent No. 5,960,792.

Other mitogenic factors, including EGF, TGF β , and FGF, have been implicated in the initiation or perpetuation of fibrosis. It may therefore be advantageous to combine a zveg3 inhibitor with one or more inhibitors of these other
35 factors.

Antibodies are preferably administered parenterally, such as by bolus injection or infusion (intravenous, intramuscular, intraperitoneal or subcutaneous) over

the course of treatment. Antibodies are generally administered in an amount sufficient to provide a minimum circulating level of antibody throughout the treatment period of between approximately 20 µg and 1 mg/kg body weight. In this regard, it is preferred to use antibodies having a circulating half-life of at least 12 hours, preferably at least 4
5 days, more preferably up to 14-21 days. Chimeric and humanized antibodies are expected to have circulatory half-lives of up to four and up to 14-21 days, respectively. In many cases it will be preferable to administer daily doses during a hospital stay, followed by less frequent bolus injections during a period of outpatient treatment. Antibodies can also be delivered by slow-release delivery systems, pumps, and other
10 known delivery systems for continuous infusion. Dosing regimens may be varied to provide the desired circulating levels of a particular antibody based on its pharmacokinetics. Thus, doses will be calculated so that the desired circulating level of therapeutic agent is maintained. Daily doses referred to above may be administered as larger, less frequent bolus administrations to provide the recited dose averaged over the
15 term of administration.

Those skilled in the art will recognize that the same principles will guide the use of other zveg3 antagonists. The dosing regimen for a given antagonist will be determined by a number of factors including potency, pharmacokinetics, and the physicochemical nature of the antagonist. For example, non-peptidic zveg3
20 antagonists may be administered enterally.

Therapeutic polynucleotides, such as antisense polynucleotides, can be delivered to patients or test animals by way of viral delivery systems. Exemplary viruses for this purpose include adenovirus, herpesvirus, retroviruses, vaccinia virus, and adeno-associated virus (AAV). Adenovirus, a double-stranded DNA virus, is
25 currently the best studied gene transfer vector for delivery of heterologous nucleic acids. For review, see Becker et al., *Meth. Cell Biol.* 43:161-89, 1994; and Douglas and Curiel, *Science & Medicine* 4:44-53, 1997. The adenovirus system offers several advantages. Adenovirus can (i) accommodate relatively large DNA inserts; (ii) be grown to high-titer; (iii) infect a broad range of mammalian cell types; and (iv) be used
30 with many different promoters including ubiquitous, tissue specific, and regulatable promoters. Because adenoviruses are stable in the bloodstream, they can be administered by intravenous injection.

By deleting portions of the adenovirus genome, larger inserts (up to 7 kb) of heterologous DNA can be accommodated. These inserts can be incorporated
35 into the viral DNA by direct ligation or by homologous recombination with a co-transfected plasmid. When intravenously administered to intact animals, adenovirus primarily targets the liver. If the adenoviral delivery system has an E1 gene deletion,

the virus cannot replicate in the host cells. However, the host's tissue (e.g., liver) will express and process (and, if a signal sequence is present, secrete) the heterologous protein.

An alternative method of gene delivery comprises removing cells from the body and introducing a vector into the cells as a naked DNA plasmid. The transformed cells are then re-implanted in the body. Naked DNA vectors are introduced into host cells by methods known in the art, including transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun, or use of a DNA vector transporter. See, Wu et al., *J. Biol. Chem.* 263:14621-14624, 1988; Wu et al., *J. Biol. Chem.* 267:963-967, 1992; and Johnston and Tang, *Meth. Cell Biol.* 43:353-365, 1994.

Activity of zveg3 antagonists can be measured *in vitro* using assays (including cell-based assays) designed to measure zveg3 activity. Antagonists will reduce the effects of zveg3 within the assay. Ligand-receptor binding can be assayed by a variety of methods well known in the art, including receptor competition assays (Bowen-Pope and Ross, *Methods Enzymol.* 109:69-100, 1985) and through the use of soluble receptors, including receptors produced as IgG fusion proteins (U.S. Patent No. 5,750,375). Receptor binding assays can be performed on cell lines that contain known cell-surface receptors for evaluation. The receptors can be naturally present in the cell, or can be recombinant receptors expressed by genetically engineered cells. Mitogenic activity can be measured using known assays, including ³H-thymidine incorporation assays (as disclosed by, e.g., Raines and Ross, *Methods Enzymol.* 109:749-773, 1985 and Wahl et al., *Mol. Cell Biol.* 8:5016-5025, 1988), dye incorporation assays (as disclosed by, for example, Mosman, *J. Immunol. Meth.* 65:55-63, 1983 and Raz et al., *Acta Trop.* 68:139-147, 1997) or cell counts. Suitable mitogenesis assays measure incorporation of ³H-thymidine into (1) 20% confluent cultures to look for the ability of zveg3 proteins to further stimulate proliferating cells, and (2) quiescent cells held at confluence for 48 hours to look for the ability of zveg3 proteins to overcome contact-induced growth inhibition. Suitable dye incorporation assays include measurement of the incorporation of the dye Alamar blue (Raz et al., *ibid.*) into target cells. See also, Gospodarowicz et al., *J. Cell. Biol.* 70:395-405, 1976; Ewton and Florini, *Endocrinol.* 106:577-583, 1980; and Gospodarowicz et al., *Proc. Natl. Acad. Sci. USA* 86:7311-7315, 1989.

The biological activities of zveg3 antagonists can be studied in non-human animals by administration of exogenous compounds, by expression of zveg3 antisense polynucleotides, and by suppression of endogenous zveg3 expression through knock-out techniques. Viral delivery systems (disclosed above) can be

employed. Zveg3 antagonists can be administered or expressed individually, in combination with other zveg3 antagonists, or in combination other compounds, including other growth factor antagonists. Test animals are monitored for changes in such parameters as clinical signs, body weight, blood cell counts, clinical chemistry, histopathology, and the like.

Effects of zveg3 antagonists on liver and kidney fibrosis can be tested in known animal models, such as the db/db mouse model disclosed by Cohen et al., *Diabetologia* 39:270-274, 1996 and Cohen et al., *J. Clin. Invest.* 95:2338-2345, 1995, transgenic animal models (Imai et al., *Contrib. Nephrol.* 107:205-215, 1994), and the CCl₄-induced cirrhosis model (Rojkind and Greenwel, *Adv. Vet. Sci. Comp. Med.* 37:333-355, 1993; Díaz-Gil et al., *J. Hepatol.* 30:1065-1072, 1999).

Effects on lung fibrosis can be assayed in a mouse model using bleomycin. The chemotherapy agent bleomycin is a known causative agent of pulmonary fibrosis in humans and can induce interstitial lung disease in mice, including an increase in the number of fibroblasts, enhanced collagen deposition, and dysregulated matrix remodeling. C57Bl/6 mice are administered bleomycin by osmotic minipump for 1 week. There follows a period of inflammation, with cutaneous toxicity beginning approximately 4-7 days after bleomycin administration and continuing for about a week, after which the mice appear to regain health. About 3-4 weeks after the finish of bleomycin delivery, the mice are sacrificed, and the lungs are examined histologically for signs of fibrosis. Scoring is based on the extent of lung fibrotic lesions and their severity. Serum is assayed for lactic dehydrogenase, an intracellular enzyme that is released into the circulation upon general cell death or injury. Lung tissue is assayed for hydroxyproline as a measure of collagen deposition.

The invention is further illustrated by the following non-limiting examples.

EXAMPLES

Example 1

A human salivary gland library was screened for a full-length clone of zveg3 by PCR. This library was an arrayed library representing 9.6×10^5 clones made in the vector pZP5x. The vector pZP5x is the same as vector pZP-9 (deposited with American Type Culture Collection, 10801 University Blvd., Manassas, VA under Accession Number 98668), but contains a cytomegalovirus promoter instead of a metallothionein promoter between the Asp718 and BamHI sites. The plasmid thus comprises a dihydrofolate reductase gene under control of the SV40 early promoter and SV40 polyadenylation site, and a cloning site to insert the gene of interest under control

of the CMV promoter and the human growth hormone (hGH) gene polyadenylation site. The working plate containing 80 pools of 12,000 colonies each was screened by PCR using oligonucleotide primers ZC19,045 (SEQ ID NO:6) and ZC19,047 (SEQ ID NO:7) with an annealing temperature of 60°C for 35 cycles. There were two strong positives, pools 58 (T-8 F1-F12) and 77 (T-7 H1-H12). The corresponding pools in the transfer plate were then screened by PCR using the same conditions. Two positives were obtained at the transfer level. The positives were T-7 H11 and T-8 F10. 5' RACE reactions were done on the transfer plate pools, and the fragments were sequenced to check zveg3 sequence and determine if a full-length clone was present. For PCR, oligonucleotide primers ZC12,700 (SEQ ID NO:8) and ZC19,045 (SEQ ID NO:6) were used at an annealing temperature of 61°C for 5 cycles, then 55°C for 30 cycles. Sequencing showed that the pool T-7 H11 had a frameshift. Transfer plate 8 pool F10 sequence appeared to be correct, so this pool of DNA was used in filter lifts.

Pool F10 from transfer plate 8 was plated and filter lifted using nylon membranes (Hybond-NTM; Amersham Corporation). Approximately 1200 colonies per plate on each of 5 filters were lifted for a total of approximately 6000 colonies. The filters were marked with a hot needle for orientation, then denatured for 6 minutes in 0.5 M NaOH and 1.5 M Tris-HCl, pH 7.2. The filters were then neutralized in 1.5 M NaCl and 0.5 M Tris-HCl, pH 7.2 for 6 minutes. The DNA was affixed to the filters using a UV crosslinker (Stratalinker®, Stratagene, La Jolla, CA) at 1200 joules. The filters were prewashed at 65°C in prewash buffer consisting of 0.25X SSC, 0.25% SDS, and 1mM EDTA. The solution was changed a total of three times over a 45-minute period to remove cell debris. Filters were prehybridized for approximately 3 hours at 65°C in 25 ml of ExpressHybTM. The probe was generated using an approximately 400-bp fragment produced by digestion of the first proprietary database clone with EcoRI and BglII and gel-purified using a spin column as disclosed above. The probe was radioactively labeled with ³²P by random priming as disclosed above and purified using a push column. ExpressHybTM solution was used for the hybridizing solution for the filters. Hybridization took place overnight at 65°C. Blots were rinsed 2X in 65°C solution 1 (2X SSC, 0.1% SDS), then washed 4 times in solution 1 at 65°C. The filters were exposed to film overnight at -80°C. There were 14 positives on the filters. 85 clones were picked from the positive areas and screened by PCR using oligonucleotide primers ZC19,045 (SEQ ID NO:6) and ZC19,047 (SEQ ID NO:7) and an annealing temperature of 60°C. Thirteen positives were obtained and streaked out for individual clones. Twenty-four colonies were picked and checked by PCR as previously described. Six positives were obtained, two of which were sequenced. Both sequences were the same and full length. The sequence is shown in SEQ ID NO:1.

A PCR panel was screened for mouse *zveg3* DNA. The panel contained 8 cDNA samples from brain, bone marrow, 15-day embryo, testis, salivary gland, placenta, 15-day embryo (Clontech Laboratories), and 17-day embryo (Clontech Laboratories) libraries.

PCR mixtures contained oligonucleotide primers ZC21,222 (SEQ ID NO:9) and ZC21,224 (SEQ ID NO:10). The reaction was run at an annealing temperature of 66°C with an extension time of 2 minutes for a total of 35 cycles using Ex Taq™ DNA polymerase (PanVera, Madison, WI) plus antibody. DNA samples found to be positive for *zveg3* by PCR and confirmed by sequencing included mouse 15-day embryo library total pool cDNA, mouse 15-day embryo (Clontech Laboratories) and 17-day embryo (both obtained from Clontech Laboratories), mouse salivary gland library total pool cDNA, and mouse testis library total pool cDNA. Fragments of about 600 bp from each of the mouse 15-day embryo library total pool cDNA, mouse 15-day embryo mcDNA, and mouse 17-day embryo mcDNA PCR products were sequenced. Sequence from the mouse 17-day embryo mcDNA and mouse 15-day embryo library total pool cDNA products confirmed the fragments to be mouse *zveg3* DNA.

The mouse 15-day embryo library was screened for full-length *zveg*f3 DNA. This library was an arrayed library representing 9.6×10^5 clones in the pCMV•SPORT 2 vector (Life Technologies, Gaithersburg, MD). The working plate, containing 80 pools of 12,000 colonies each, was screened by PCR using oligonucleotide primers ZC21,223 (SEQ ID NO:11) and ZC21,224 (SEQ ID NO:10) with an annealing temperature of 66°C for 35 cycles. Eighteen positives were obtained. Fragments from four pools (A2, A10, B2, and C4) were sequenced; all were confirmed to encode *zveg*f3. Additional rounds of screening using the same reaction conditions and pools from the working and source plates identified one positive pool (5D).

Positive colonies were screened by hybridization. Pool 5D from original source plate #5 was plated at about 250 colonies per plate and transferred to nylon membranes (Hybond-NTM; Amersham Corporation, Arlington Heights, IL). Five filters were lifted for a total of ~1250 colonies. The filters were marked with a hot needle for orientation, then denatured for 6 minutes in 0.5 M NaOH and 1.5 M Tris-HCl, pH 7.2. The filters were then neutralized in 1.5 M NaCl and 0.5 M Tris-HCl, pH 7.2 for 6 minutes. The DNA was fixed to the filters using a UV crosslinker (Stratalinker®, Stratagene, La Jolla, CA) at 1200 joules. A probe was generated by PCR using oligonucleotide primers ZC21,223 (SEQ ID NO:11) and ZC21,224 (SEQ ID NO:10), and a mouse 15-day embryo template at an annealing temperature of 66°C for 35

cycles. The PCR fragment was gel purified using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA). The DNA was radioactively labeled with ³²P using a commercially available kit (Rediprime™ II random-prime labeling system; Amersham Corp., Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a commercially available push column (NucTrap® column; Stratagene, La Jolla, CA; see U.S. Patent No. 5,336,412). The filters were prewashed at 65°C in prewash buffer consisting of 0.25X SSC, 0.25% SDS and 1mM EDTA. The solution was changed a total of three times over a 45-minute period to remove cell debris. Filters were prehybridized overnight at 65°C in 25 ml of a hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc., Palo Alto, CA), then hybridized overnight at 65°C in the same solution. Filters were rinsed twice at 65°C in pre-wash buffer (0.25X SSC, 0.25% SDS, and 1mM EDTA), then washed twice in pre-wash buffer at 65°C. Filters were exposed to film for 2 days at -80°C. There were 10 positives on the filters. 3 clones were picked from the positive areas, streaked out, and 15 individual colonies from these three positives were screened by PCR using primers ZC21,223 (SEQ ID NO:11) and ZC21,334 (SEQ ID NO:12) at an annealing temp of 66°C. Two positives were recovered and sequenced. Both sequences were found to be the same and encoded full-length mouse zveg3 (SEQ ID NO:4).

The amino acid sequence is highly conserved between mouse and human zveg3s, with an overall amino acid sequence identity of 87%. The secretory peptide, CUB domain, inter-domain, and growth factor domain have 82%, 92%, 79% and 94% amino acid identity, respectively.

25 Example 3

A mammalian cell expression vector for the growth factor domain of zveg3 was constructed by joining the zveg3 fragment to a sequence encoding an optimized t-PA secretory signal sequence (U.S. Patent No. 5,641,655) in the linearized pZMP11 vector downstream of the CMV promoter. The plasmid pZMP11 is a mammalian expression vector containing an expression cassette having the CMV immediate early promoter, a consensus intron from the variable region of mouse immunoglobulin heavy chain locus, Kozak sequences, multiple restriction sites for insertion of coding sequences, a stop codon, and a human growth hormone terminator. The plasmid also contains an IRES element from poliovirus, the extracellular domain of CD8 truncated at the C-terminal end of the transmembrane domain, an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, enhancer and origin of replication, a DHFR gene, the SV40 terminator, and

the URA3 and CEN-ARS sequences required for selection and replication in *S. cerevisiae*. The resulting vector was designated pZMP11/zv3GF-otPA.

BHK 570 cells were transfected with pZMP11/zv3GF-otPA by liposome-mediated transfection (using LipofectamineTM; Life Technologies) and
5 cultured according to conventional procedures.

BHK cell-conditioned media was adjusted to 20 mM MES at pH 5.5. A column of cation exchange resin (Poros® HS 50; PerSeptive Biosystems, Framingham, MA) (2-cm diameter; 50 ml bed volume) was equilibrated in 20 mM MES, 150 mM NaCl, pH 5.5. The adjusted media was pumped into the column at 20 ml/minute.
10 When loading was completed, the column was washed, first in 20 mM MES, 150 mM NaCl, pH 5.5, then with the same composition buffer at pH 6.0. Once the baseline was back to zero absorbance, the column was eluted with a 10 column volume gradient to 20mM MES, 1M NaCl, pH 6.0. The zveg3 growth factor domain eluted at 25 to 50 mS conductivity during the evolving gradient. Reducing SDS-PAGE revealed a
15 distinct band at 20 kD, which was confirmed as zveg3 by Western blotting. This material was pooled and prepared for loading to a tandem column array comprising a strong anion exchange resin (Poros® HQ 50; PerSeptive Biosystems) followed by an immobilized metal (nickel) affinity column in series. The system of columns was equilibrated in 20 mM MOPS buffer at pH 7.0. The vegf3 pool was in-line diluted at
20 1:10 (V:V) with the MOPS equilibration buffer while loading. After loading was completed the column series was washed with 20 mM MOPS pH 7.0 buffer until baseline absorbance was obtained. The nickel column was then disconnected from the anion exchanger and washed with 20 mM MOPS pH 7.0 containing 150 mM NaCl. The columnwais then eluted with a 1 column volume gradient between the last washing
25 buffer and the same buffer containing 20 mM imidazole at pH 7.0. The fractions containing the zveg3 growth factor domain were pooled and concentrated using 5 kD cutoff membrane in preparation for buffer exchange and polishing on a size exclusion column equilibrated in PBS.

30 Example 4

To make transgenic animals expressing zveg3 genes requires adult, fertile males (studs) (B6C3f1, 2-8 months of age (Taconic Farms, Germantown, NY)), vasectomized males (duds) (B6D2f1, 2-8 months, (Taconic Farms)), prepubescent fertile females (donors) (B6C3f1, 4-5 weeks, (Taconic Farms)) and adult fertile females
35 (recipients) (B6D2f1, 2-4 months, (Taconic Farms)).

The donors are acclimated for 1 week, then injected with approximately 8 IU/mouse of Pregnant Mare's Serum gonadotrophin (Sigma, St. Louis, MO) I.P., and

46-47 hours later, 8 IU/mouse of human Chorionic Gonadotropin (hCG (Sigma)) I.P. to induce superovulation. Donors are mated with studs subsequent to hormone injections. Ovulation generally occurs within 13 hours of hCG injection. Copulation is confirmed by the presence of a vaginal plug the morning following mating.

- 5 Fertilized eggs are collected under a surgical scope (Leica MZ12 Stereo Microscope, Leica, Wetzlar, Germany). The oviducts are collected and eggs are released into urinalysis slides containing hyaluronidase (Sigma Chemical Co.). Eggs are washed once in hyaluronidase, and twice in Whitten's W640 medium (Table 2; all reagents available from Sigma Chemical Co.) that has been incubated with 5% CO₂,
 10 5% O₂, and 90% N₂ at 37°C. The eggs are stored in a 37°C/5% CO₂ incubator until microinjection.

Table 2

	<u>mgs/200 ml</u>	<u>mgs/500 ml</u>
NaCl	1280	3200
KCl	72	180
KH ₂ PO ₄	32	80
MgSO ₄ · 7H ₂ O	60	150
Glucose	200	500
Ca ²⁺ Lactate	106	265
Benzylpenicillin	15	37.5
Streptomycin SO ₄	10	25
NaHCO ₃	380	950
Na Pyruvate	5	12.5
H ₂ O	200 ml	500 ml
500 mM EDTA	100 µl	250 µl
5% Phenol Red	200 µl	500 µl
BSA	600	1500

- 15 Zveg3 cDNA is inserted into the expression vector pHB12-8. Vector pHB12-8 was derived from p2999B4 (Palmiter et al., *Mol. Cell Biol.* 13:5266-5275, 1993) by insertion of a rat insulin II intron (ca. 200 bp) and polylinker (Fse I/Pme I/Asc I) into the Nru I site. The vector comprises a mouse metallothionein (MT-1) promoter (ca. 750 bp) and human growth hormone (hGH) untranslated region and

polyadenylation signal (ca. 650 bp) flanked by 10 kb of MT-1 5' flanking sequence and 7 kb of MT-1 3' flanking sequence. The cDNA is inserted between the insulin II and hGH sequences.

10-20 micrograms of plasmid DNA is linearized, gel-purified, and
 5 resuspended in 10 mM Tris pH 7.4, 0.25 mM EDTA pH 8.0, at a final concentration of 5-10 nanograms per microliter for microinjection.

Plasmid DNA is microinjected into harvested eggs contained in a drop of W640 medium overlaid by warm, CO₂-equilibrated mineral oil. The DNA is drawn
 10 into an injection needle (pulled from a 0.75mm ID, 1mm OD borosilicate glass capillary) and injected into individual eggs. Each egg is penetrated with the injection needle into one or both of the haploid pronuclei.

Picoliters of DNA are injected into the pronuclei, and the injection
 needle is withdrawn without coming into contact with the nucleoli. The procedure is repeated until all the eggs are injected. Successfully microinjected eggs are transferred
 15 into an organ tissue-culture dish with pregassed W640 medium for storage overnight in a 37°C/5% CO₂ incubator.

The following day, 2-cell embryos are transferred into pseudopregnant recipients. The recipients are identified by the presence of copulation plugs, after copulating with vasectomized duds. Recipients are anesthetized and shaved on the
 20 dorsal left side and transferred to a surgical microscope. A small incision is made in the skin and through the muscle wall in the middle of the abdominal area outlined by the ribcage, the saddle, and the hind leg, midway between knee and spleen. The reproductive organs are exteriorized onto a small surgical drape. The fat pad is stretched out over the surgical drape, and a baby serrefine (Roboz, Rockville, MD) is
 25 attached to the fat pad and left hanging over the back of the mouse, preventing the organs from sliding back in.

With a fine transfer pipette containing mineral oil followed by alternating W640 and air bubbles, 12-17 healthy 2-cell embryos from the previous day's
 30 injection are transferred into the recipient. The swollen ampulla is located and holding the oviduct between the ampulla and the bursa, and a nick in the oviduct is made with a 28 g needle close to the bursa, making sure not to tear the ampulla or the bursa.

The pipette is transferred into the nick in the oviduct, and the embryos
 are blown in, allowing the first air bubble to escape the pipette. The fat pad is gently pushed into the peritoneum, and the reproductive organs are allowed to slide in. The
 35 peritoneal wall is closed with one suture, and the skin is closed with a wound clip. The mice recuperate on a 37°C slide warmer for a minimum of 4 hours.

006924 102200

35 Analysis of the mRNA expression level of each transgene is done using
an RNA solution hybridization assay or real-time PCR on an ABI Prism 7700 (PE
Applied Biosystems, Inc., Foster City, CA) following the manufacturer's instructions.

5

10

35

expression with low body weight, low hematocrit, and low platelet count. One AEO/zveg3 transgenic mouse (male) with a low level of expression exhibited liver sinusoidal cell proliferation.

Histological analysis was carried out on livers from N1 transgenic mice overexpressing full-length zveg3 from the AEO promoter and nontransgenic littermate controls at 8, 16, 22, and 33 weeks of age. Similar changes were seen in male and female animals. H & E and trichrome stains indicate a definite increase in number of liver sinusoidal (stellate) cells and increased perisinusoidal extracellular matrix (EMC) deposition at 8 weeks of age. There was a persistent increase in number of stellate cells and in the amount and thickness of perisinusoidal EMC as well as some perivenular EMC deposition by 16 weeks. At 22 weeks similar changes were seen, but with an increase in incidence and severity of perivenular fibrosis (steato fibrosis). Changes similar to those at 16 and 22 weeks were observed at 33 weeks, however some animals had fibrotic banding and multiple, encapsulated areas in which hepatocytes appeared enlarged and vacuolated. These areas tended to be surrounded by a fibrous tissue capsule, and the hepatocytes within these areas appeared normal. These changes were consistent with early cirrhosis.

Similar changes were seen in 8-week-old N2 mice and in breeder males sacrificed at approximately 32-34 weeks.

Example 5

For construction of adenovirus vectors, the protein coding region of human zveg3 was amplified by PCR using primers that added PmeI and AscI restriction sites at the 5' and 3' termini respectively. PCR primers ZC20,180 (SEQ ID NO:17) and ZC20,181 (SEQ ID NO:18) were used with a full-length zveg3 cDNA template in a PCR reaction as follows: incubation at 95°C for 5 minutes; followed by 15 cycles at 95°C for 1 min., 61°C for 1 min., and 72°C for 1.5 min.; followed by 72°C for 7 min.; followed by a 4°C soak. The reaction product was loaded onto a 1.2 % low-melting-temperature agarose gel in TAE buffer (0.04 M Tris-acetate, 0.001 M EDTA). The zveg3 PCR product was excised from the gel and purified using a commercially available kit comprising a silica gel membrane spin column (QIAquick™ PCR Purification Kit and gel cleanup kit; Qiagen, Inc.) as per kit instructions. The zveg3 product was then digested with PmeI and AscI, phenol/chloroform extracted, EtOH precipitated, and rehydrated in 20 ml TE (Tris/EDTA pH 8). The 1038 bp zveg3 fragment was then ligated into the PmeI-AscI sites of the transgenic vector pTG12-8 (also known as pHB12-8; see Example 4) and transformed into *E. coli* DH10B™

competent cells by electroporation. Clones containing *zveg3* were identified by plasmid DNA miniprep followed by digestion with *PmeI* and *AscI*. A positive clone was sequenced to insure that there were no deletions or other anomalies in the construct. The sequence of *zveg3* cDNA was confirmed.

DNA was prepared using a commercially available kit (Maxi Kit, Qiagen, Inc.), and the 1038bp zveg3 cDNA was released from the pTG12-8 vector using PmeI and AscI enzymes. The cDNA was isolated on a 1% low melting temperature agarose gel and was excised from the gel. The gel slice was melted at 70°C, and the DNA was extracted twice with an equal volume of Tris-buffered phenol and precipitated with EtOH. The DNA was resuspended in 10 µl H₂O.

The zvegf3 cDNA was cloned into the EcoRV-AscI sites of a modified pAdTrack-CMV (He, T-C. et al., *Proc. Natl. Acad. Sci. USA* 95:2509-2514, 1998). This construct contains the green fluorescent protein (GFP) marker gene. The CMV promoter driving GFP expression was replaced with the SV40 promoter, and the SV40 polyadenylation signal was replaced with the human growth hormone polyadenylation signal. In addition, the native polylinker was replaced with FseI, EcoRV, and AscI sites. This modified form of pAdTrack-CMV was named pZyTrack. Ligation was performed using a commercially available DNA ligation and screening kit (Fast-Link™ kit; Epicentre Technologies, Madison, WI). Clones containing zvegf3 were identified by digestion of mini prep DNA with FseI and AscI. In order to linearize the plasmid, approximately 5 µg of the resulting pZyTrack zvegf3 plasmid was digested with PmeI. Approximately 1 µg of the linearized plasmid was cotransformed with 200 ng of supercoiled pAdEasy (He et al., *ibid.*) into *E. coli* BJ5183 cells (He et al., *ibid.*). The co-transformation was done using a Bio-Rad Gene Pulser at 2.5 kV, 200 ohms and 25 µFa. The entire co-transformation mixture was plated on 4 LB plates containing 25 µg/ml kanamycin. The smallest colonies were picked and expanded in LB/kanamycin, and recombinant adenovirus DNA was identified by standard DNA miniprep procedures. Digestion of the recombinant adenovirus DNA with FseI and AscI confirmed the presence of the zvegf3 insert. The recombinant adenovirus miniprep DNA was transformed into *E. coli* DH10B™ competent cells, and DNA was prepared using a Maxi Kit (Qiagen, Inc.) according to kit instructions.

Approximately 5 µg of recombinant adenoviral DNA was digested with PacI enzyme (New England Biolabs) for 3 hours at 37°C in a reaction volume of 100 µl containing 20-30U of PacI. The digested *DNA* was extracted twice with an equal volume of phenol/chloroform and precipitated with ethanol. The DNA pellet was resuspended in 10µl distilled water. A T25 flask of QBI-293A cells (Quantum Biotechnologies, Inc. Montreal, Qc. Canada), inoculated the day before and grown to

Seven days after transfection of 293A cells with the recombinant adenoviral DNA, the cells expressed GFP and started to form foci. The crude viral lysate was collected using a cell scraper to *collect* all of the 293A cells. The lysate was transferred to a 50-ml conical tube. To release most of the virus particles from the cells, three freeze/thaw cycles were done in a dry ice/ethanol bath and a 37° waterbath.

Secondary (2°) amplification of zveg3 rAdV was obtained as follows: Twenty 15-cm tissue culture dishes of 293A cells were prepared so that the cells were 80-90% confluent. All but 20 ml of 5% MEM media was removed, and each dish was inoculated with 300-500 µl of the 1° amplified rAdv lysate. After 48 hours the cells were lysed from virus production, the lysate was collected into 250-ml polypropylene centrifuge bottles, and the rAdV was purified.

NP-40 detergent was added to a final concentration of 0.5% to the
35 bottles of crude lysate in order to lyse all cells. Bottles were placed on a rotating
platform for 10 minutes agitating as fast as possible without the bottles falling over.
The debris was pelleted by centrifugation at 20,000 X G for 15 minutes. The

supernatant was transferred to 250-ml polycarbonate centrifuge bottles, and 0.5 volume of 20% PEG8000/2.5 M NaCl solution was added. The bottles were shaken overnight on ice. The bottles were centrifuged at 20,000 X G for 15 minutes and, the supernatant was discarded into a bleach solution. Using a sterile cell scraper, the white, virus/PEG precipitate from 2 bottles was resuspended in 2.5 ml PBS. The resulting virus solution was placed in 2-ml microcentrifuge tubes and centrifuged at 14,000 X G in the microcentrifuge for 10 minutes to remove any additional cell debris. The supernatant from the 2-ml microcentrifuge tubes was transferred into a 15-ml polypropylene snapcap tube and adjusted to a density of 1.34 g/ml with CsCl. The volume of the virus solution was estimated, and 0.55 g/ml of CsCl was added. The CsCl was dissolved, and 1 ml of this solution weighed 1.34 g. The solution was transferred to 3.2-ml, polycarbonate, thick-walled centrifuge tubes and spun at 348,000 X G for 3-4 hours at 25°C. The virus formed a white band. Using wide-bore pipette tips, the virus band was collected.

The virus recovered from the gradient had a large amount of CsCl which had to be removed before it was used on cells. Commercially available ion-exchange columns (PD-10 columns prepacked with Sephadex® G-25M; Pharmacia Biotech, Piscataway, NJ) were used to desalt the virus preparation. The column was equilibrated with 20 ml of PBS. The virus was loaded and allowed to run into the column. 5 ml of PBS was added to the column, and fractions of 8-10 drops were collected. The optical density of a 1:50 dilution of each fraction was determined at 260 nm on a spectrophotometer. A clear absorbance peak was present between fractions 7-12. Peak fractions were pooled, and the optical density (OD) of a 1:25 dilution was determined. OD was converted to virus concentration using the formula: $(OD \text{ at } 260\text{nm})(25)(1.1 \times 10^{12}) = \text{virions/ml}$. The OD of a 1:25 dilution of the zvegf3 rAdV was 0.145, giving a virus concentration of 4×10^{12} virions/ml.

To store the virus, glycerol was added to the purified virus to a final concentration of 15%, mixed gently but effectively, and stored in aliquots at -80°C.

A protocol developed by Quantum Biotechnologies, Inc. (Montreal, Canada) was followed to measure recombinant virus infectivity. Briefly, two 96-well tissue culture plates were seeded with 1×10^4 293A cells per well in MEM containing 2% fetal bovine serum for each recombinant virus to be assayed. After 24 hours 10-fold dilutions of each virus from 1×10^{-2} to 1×10^{-14} were made in MEM containing 2% fetal bovine serum. 100 µl of each dilution was placed in each of 20 wells. After 5 days at 37°C, wells were read either positive or negative for Cytopathic Effect (CPE), and a value for plaque forming units (pfu)/ml was calculated.

5 To calculate titer of the undiluted virus sample: the factor, “F” = 1+d(S-0.5); where “S” is the sum of the ratios (R); and “d” is Log10 of the dilution series, for example, “d” is equal to 1 for a ten-fold dilution series. The titer of the undiluted sample is $T = 10^{(1+F)} = \text{TCID}_{50}/\text{ml}$. To convert $\text{TCID}_{50}/\text{ml}$ to pfu/ml, 0.7 is subtracted from the exponent in the calculation for titer (T).

10

Example 6

15

20

25

30

35

The huzveg3 peptide-specific antibodies were affinity purified from the rabbit serum using an CNBr-Sepharose® 4B peptide column (Pharmacia Biotech) that was prepared using 10 mg of the respective peptides per gram CNBr-Sepharose®, followed by dialysis in PBS overnight. Peptide specific-huzveg3 antibodies were characterized by an ELISA titer check using 1 µg/ml of the appropriate peptide as an antibody target. The huzveg3-1 peptide-specific antibodies had a lower limit of detection (LLD) of 500 pg/ml by ELISA on the appropriate antibody target and recognize full-length recombinant protein (MBP-fusion) by ELISA. The huzveg3-2 peptide-specific antibodies had an LLD of 1 ng/ml by ELISA. The huzveg3-3 peptide-specific antibodies had an LLD of 50 pg/ml by ELISA and recognized recombinant protein by Western Blot analysis. The huzveg3-4 peptide-specific antibodies had an LLD of 50 pg/ml by ELISA and recognized recombinant protein by Western Blot analysis.

15 Example 8

Mouse hybridomas producing monoclonal antibodies (MAbs) specific for recombinant human zveg3 growth factor domain (GFD) protein were generated using purified, untagged, recombinant human zveg3 GFD produced in BHK cells (huzveg3-GFD-BHK). Ten BALB/c mice were each injected IP on day 1 with 20 µg of huzveg3-GFD-BHK mixed 1:1 (v/v) in complete Freund's adjuvant. Each mouse was subsequently injected IP with 10 µg of huzveg3-GFD-BHK mixed 1:1 in incomplete Freund's adjuvant on days 15, 29, 41, 57, 71, 89 and 115. On day 118, splenocytes and lymphocytes from enlarged lymph nodes of two mice with the highest anti-huzveg3 antibody titer (as determined in a biotinylated huzveg3-GFD capture ELISA; see below) were fused at a 2.76:1 ratio with the X63-Ag8.653 mouse myeloma cell line (Kearney et al., *J. Immunol.* 123:1548-1550, 1979) essentially as disclosed by Lane (*J. Immunol. Methods* 81:223-228, 1985). The fusion mixture was plated into 24 96-well plates at an average density of 1.2×10^5 total cells/well in Iscove's modified Dulbecco's medium (IMDM; Life Technologies, Inc., Gaithersburg, MD) containing 10% fetal clone I serum (HyClone Laboratories, Inc., Logan, UT), 10% hybridoma cloning supplement (BM Condimed® H1; Roche Diagnostics Corp., Indianapolis, IN), 2 mM L-glutamine (Life Technologies, Inc.), 100 U/mL penicillin G sodium (Life Technologies, Inc.), and 100 µg/mL streptomycin sulfate (Life Technologies, Inc.). Wells were fed on days 4 and 7 by aspiration and replacement of approximately three-fourths of the media contents in each well. This fusion was designated HH1.

Anti-huzveg3 mAbs of the IgG class were detected on days 9/10 post-fusion using a biotinylated huveg3-GFD capture ELISA. Wells of plates (Immulon®

II; Dynex Technologies, Chantilly, VA) were coated with 1 µg/mL of goat anti-mouse IgG (obtained from Kirkegaard & Perry Laboratories, Gaithersburg, MD) in 0.05 M carbonate/bicarbonate buffer (Sigma, St. Louis, MO), 50 µL/well. Plates were incubated at 4°C overnight, then washed three times with phosphate buffered saline containing 0.05% polyoxyethylenesorbitan monolaurate (Tween 20) (PBST) to remove unbound antibody. Wells were blocked with PBST containing 1% (w/v) bovine serum albumin (Sigma) (PTB buffer), 200 µL/well, for 1 hour at room temperature (RT), then washed as above. Culture supernatants from the fusion plates were replica-plated onto antibody-coated plates, 100 µL/well, and incubated at RT for 1 hour. Human zveg3-GFD protein was biotinylated with sulfo-NHS-LC-biotin (Pierce Chemical Company, Rockford, IL) according to the manufacturer's instructions for 45 minutes at RT. The reaction was stopped with 2M glycine. Biotinylated protein was diluted to 1 µg/mL in PTB buffer. The plates were washed four times with PBST, biotinylated huzveg3-GFD was added at 100 µL/well, and the plates were incubated at RT for 1 hour. The plates were then washed four times with PBST, and HRP-conjugated streptavidin (Pierce Chemical Company) was diluted to 0.5 µg/mL in PTB buffer and added 100 to the plates at µL/well. The plates were incubated at RT for 1 hour, then washed five times with PBST. TMB substrate (EIA chromagen, Cat. # R6R; Genetic Systems Corp.) was diluted 1:100 in substrate buffer (Genetic Systems Corp.) and added to the plates at 100 µL/well. The plates were incubated 10-15 minutes at RT in the dark. Color development was stopped by the addition of 100 µL/well of 1N H₂SO₄. Optical density was read on an ELISA plate reader (Molecular Devices, Sunnyvale, CA) at wavelengths 450 nm (L1) and 650 nm (L2). Final OD measurement was determined by the formula L1-L2.

78 master wells contained antibody that was capable of capturing biotinylated huzveg3-GFD. These master wells were expanded for hybridoma cryopreservation and additional supernatant generation. ELISA analysis of the expanded master well supernatants demonstrated that 27 of the original 78 master wells retained antibody specific for huzveg3-GFD, with 20 of 27 possessing significant reactivity with the antigen.

Six of the designated master wells were cloned by limiting dilution at a density of less than one cell per well. Monoclonality was assessed by microscopic examination of wells for a single focus of cell growth. Clones were tested for specific antibody by the same assay used in the master screen. Five to six of the strongest clones were briefly expanded. Supernatants from these clones was then serially diluted and tested by ELISA to identify the three best antibody-producing clones. Upon

verification, the best clone from each master well was subsequently recloned and assayed as described above.

Six of the highest titered second round clones were adapted to growth in fusion/cloning medium minus the addition of hybridoma cloning supplement. Each of the clones was subsequently adapted to growth in a production medium formulation consisting essentially of Dulbecco's modified Eagle's medium + 2.5% fetal clone I serum and various supplements. Supernatants were again serially diluted and tested by ELISA on huzveg3-GFD to identify the highest titered and next highest titered clones (designated the primary and secondary clones, respectively). MAb produced by each set of clones was then evaluated for IgG subclass using the Mouse Hybridoma Subtyping Kit (Cat. #1183117; Roche Diagnostics Corp.). Clones HH1-24, -40, -57 and -76 were all found to produce an IgG₁ antibody. Clones HH1-58 and -78 produced an IgG_{2b} antibody. All antibodies possessed a κ light chain.

15 Example 9

Recombinant zveg3 was analyzed for mitogenic activity on rat stellate cells (Greenwel et al., *Laboratory Invest.* 65:644, 1991; Greenwel et al., *Laboratory Invest.* 69:210, 1993). Stellate cells were plated at a density of 2,000 cells/well in 96-well culture plates and grown for approximately 72 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for 20 hours in serum-free DMEM/Ham's F-12 medium containing insulin (5 μ g/ml), transferrin (20 μ g/ml), and selenium (16 pg/ml) (ITS). At the time of the assay, the medium was removed, and test samples were added to the wells in triplicate. Test samples consisted of either conditioned media (CM) from adenovirally-infected HaCaT human keratinocyte cells (Boukamp et al., *J. Cell. Biol.* 106:761-771, 1988) expressing full-length zveg3, purified growth factor domain expressed in BHK cells, or control media from cells infected with parental adenovirus (Zpar) containing an expression unit for green fluorescent protein. The CM was concentrated 10-fold using a 15-ml centrifugal filter device with a 10K membrane filter (Ultrafree®; Millipore Corp., Bedford, MA), then diluted back to 3x with ITS medium and added to the cells. Purified protein in a buffer containing 0.1% BSA was serially diluted into ITS medium at concentrations of 1 μ g/ml to 1 ng/ml and added to the test plate. A control buffer of 0.1% BSA was diluted identically to the highest concentration of zveg3 protein and added to the plate. For measurement of [³H]thymidine incorporation, 20 μ l of a 50 μ Ci/ml stock in DMEM was added directly to the cells, for a final activity of 1 μ Ci/well. After another 24-hour incubation, mitogenic activity was assessed by measuring the uptake of [³H]thymidine. Media were removed, and cells were incubated with 0.1 ml of trypsin

until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.) and counted on a microplate scintillation counter (Topcount®; Packard Instrument Co.).

Results, presented in Table 3, demonstrated that zveg3 CM had approximately 4.4-fold higher mitogenic activity on stellate cells over control CM, and purified protein at 100 ng/ml caused a maximal 14-fold increase in [³H]thymidine incorporation over the buffer control.

Table 3

<u>Sample</u>	<u>CPM Incorporated</u>	
	<u>Mean</u>	<u>St. dev.</u>
zveg3 (2x CM)	42489	1306
Zpar (2x CM)	9629	540
zveg3 GF domain, 100 ng/ml	77540	4142
zveg3 GF domain, 33.3 ng/ml	74466	18142
zveg3 GF domain, 11.1 ng/ml	52462	6239
zveg3 GF domain, 3.7 ng/ml	15128	4989
Buffer control	5618	573
PDGF-BB 20 ng/ml	19741	2075
PDGF-AA 20 ng/ml	33133	3325
Media alone (basal response)	6765	226

Example 10

Northern blot analysis was performed on 2 µg samples of poly(A)⁺ RNA from five mouse prostate cell lines (designated Jakotay, Nelix, Paris, Torres, and Tuvak). Total RNA from mouse and rat liver were used as controls (20 µg each). An approximately 680-bp DNA probe was generated by PCR using oligonucleotide primers ZC21,222 (SEQ ID NO:9) and ZC21,224 (SEQ ID NO:10) and a mouse zveg3 full-length cDNA clone as a template. The DNA probe was purified by conventional procedures using a commercially available kit (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA). The probe was radioactively labeled with ³²P using a commercially available kit (Rediprime™ II DNA Labeling system; Amersham,

Arlington Heights, IL) according to the manufacturer's specifications. The probe was purified using a push column (NucTrap® column; Stratagene, La Jolla, CA; see U.S. Patent No. 5,336,412). A commercially available hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories, Inc., Palo Alto, CA) was used for the blots. Hybridization took place overnight at 65°C. The blots were then washed four times in 2X SCC and 0.1% SDS at room temperature, followed by two washes in 0.1X SSC and 0.1% SDS at 55° C. One transcript size was detected at approximately 4 kb in Jakotay, Nelix, Paris and Tuvak cell lines. Signal intensity was highest for Jakotay, then Nelix and Tuvak.

Example 11

Recombinant human zveg3 GFD produced in BHK cells was assayed for growth factor activity using an osteoblast cell line containing a luciferase reporter gene construct. CCC4 cells (derived from p53 knock-out mice) transfected with the reporter construct were plated in 96-well plates (BioCoat™; Becton Dickinson, Franklin Lakes, NJ) at 1×10^4 cells/well in 1% FBS alpha MEM (JRH Biosciences, Lexena, KS) containing L-Glutamine and sodium pyruvate (Life Technologies, Inc.). The cells were cultured for 24 hours, the medium was removed, and the wells were washed twice with wash buffer (1% BSA in PBS), 100 µl/well at 37°C. Zveg3, PDGF-AA, and PDGF-BB were diluted in assay medium (alpha MEM containing 1% BSA, 10 mM Hepes, L-Glutamate, and sodium pyruvate), and 100-µl samples were added to the wells. The plates were incubated 4 hours at 37°C, then the wells were washed with wash buffer. Lysis reagent (Cell Culture Lysis Reagent; Promega Corporation, Madison, WI) was diluted 1:5 in water, and 25 µl was added to each well. The plates were incubated 10 minutes at room temperature, then transferred to -20°C. Luciferase substrate (Luciferase Assay System; Promega Corporation) was diluted 1:1 with assay medium and added to the wells. The plates were read on a luminometer. Results, shown in Table 4, indicted that all three growth factors exhibited mitogenic activity on the CCC4 cells.

Table 4

Concentration	Activity (Luminometer Units)		
	(ng/ml)	Zveg3	PDGF-BB PDGF-AA
0		3.5	3.1 3.2
0.4		5.2	7.7 8.4
0.8		5.6	10.7 9.5
1.5		7.6	14 9.4
3.1		10.6	19.9 11.4
6.2		12.5	25.1 12
12.5		14.2	26.1 12.3
25		16	25.7 11.9
50		19.4	25.8 12.6
100		20.6	23.8 14

5 Example 12

A study was undertaken to test whether adenovirally delivered zveg3 stimulated cell proliferation as determined by incorporation of bromodeoxyuridine (BrdU) into tissues.

Mice (male, C57Bl, 7 weeks old) were divided into three groups. On day 0, parental or zveg3 adenovirus was administered to the first (n=11) and second (n=12) groups, respectively, via the tail vein, with each mouse receiving a dose of $\sim 1 \times 10^{11}$ particles in ~ 0.1 ml volume. The third group (n=8) received no treatment. Each mouse was given two intraperitoneal doses of 3 mg of freshly made BrdU solution at approximately 24 and 12 hours prior to sacrifice. On days 2, 4, 6, 8, and 10, two mice from each treatment group and one or two untreated mice were sacrificed, and tissues and blood were harvested. Samples were analyzed for complete blood count (CBC) and serum chemistry, and slides were prepared for manual differential blood and marrow progenitor cell analysis. One femur, lung, heart, thymus, liver, kidney, spleen, pancreas, duodenum, and mesenteric lymph nodes were submitted for standard histology and assessment of BrdU incorporation. The lining of the duodenum served as the control tissue for BrdU incorporation.

In addition, two mice that received approximately half the dose of zveg3 adenovirus particles and one mouse that received the full dose of parental adenovirus were sacrificed and analyzed as described above on day 16.

A piece of liver from each mouse was saved for mRNA assay of adenovirus protein to examine the time course of expression of the adenovirus preparations.

Beginning on day 6, most of the animals treated with either adenovirus had visibly enlarged livers and spleens compared to the untreated mice. The livers of

the zveg3 adenovirus-treated mice tended to look more pale than animals treated with the parental virus. Proliferation of sinusoidal cells was observed in liver. Visual inspection suggested that these cells were stellate cells and/or fibroblasts. Spleen color was the same in both groups. Most of the animals that received the zveg3 adenovirus

5 had paler femur shafts, with the marrow lighter in color.

Peripheral blood CBCs showed a possible difference in platelet counts, but not in RBC or WBC counts between zveg3 and parental virus-treated animals. In comparison to the untreated and parental virus-treated groups, the zveg3 group had lower platelet counts on days 2, 4, 6, and 8, but not on day 10. The mean platelet

10 volume (average size of individual platelets) in the zveg3 group also tended to be greater, consistent with a relative increase in the larger, immature platelet population.

BrdU labeling showed increased cell proliferation in kidney, mainly in the medulla and to a lesser extent in the cortex. Proliferating cells appeared to be interstitial cells, which may have included fibroblasts and/or mesangial cells.

15

Example 13

Human zveg3 growth factor domain protein produced in BHK cells was tested for the ability to stimulate production of TGF- β in stellate cells. Rat hepatic stellate cells (obtained from Dr. Nelson Fausto, University of Washington) were plated

20 in 48-well tissue culture clusters (Costar®; Corning, Corning, NY) in DMEM growth medium (Life Technologies, Inc.) supplemented with pyruvate and 10% serum (HyClone Laboratories, Inc.). At confluence, the medium was changed to serum-free medium by substituting 0.1% BSA (Fraction V; Sigma, St. Louis, MO) for serum. 48 hours later, the medium was changed, replaced with the same serum-free medium, and

25 the zveg3 GFD protein was added at 100 ng/ml in the wells. 48 hours later, conditioned media were collected and spun down to get rid of any floating cells or debris. Total TGF- β 1 levels were determined in these media using a commercially available ELISA kit (R&D Systems, St Paul, MN). Stimulation of stellate cells with

30 100 ng/ml zveg3 GFD resulted in an approximately 5-fold increase in the production of TGF- β compared to a BSA control.

Example 14

Recombinant human zveg3 GFD was analyzed for mitogenic activity on human mesangial cells (Clonetics, San Diego, CA). Mesangial cells were plated at a

35 density of 2,000 cells/well in 96-well culture plates and grown for approximately 72 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for 20 hours in serum-free DMEM/Ham's F-12 medium containing

insulin (5 µg/ml), transferrin (20 µg/ml), and selenium (16 pg/ml) (ITS). At the time of the assay, the medium was removed, and test samples were added to the wells in triplicate. Test samples consisted of purified zveg3 GFD at 3, 10, 30 and 100 ng/ml, and PDGF-AA, PDGF-AB and PDGF-BB at 30 ng/ml in a buffer containing 0.1% BSA. Samples were serially diluted into ITS medium and added to the test plate. A control buffer of 0.1% BSA was diluted identically to the highest concentration of zveg3 protein and added to the plate. 10% fetal bovine serum (FBS) was used as a positive control. For measurement of [³H]thymidine incorporation, 20 µl of a 50 µCi/ml stock in DMEM was added directly to the cells, for a final activity of 1 µCi/well. After another 24-hour incubation, mitogenic activity was assessed by measuring the uptake of [³H]thymidine. Media were removed, and cells were incubated with 0.1 ml of trypsin until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.), and counted on a microplate scintillation counter (Topcount®; Packard Instrument Co.). Results, shown in Table 5, demonstrated that zveg3 GFD had approximately 3-fold higher mitogenic activity cells over PDGF-AA and comparable activity to PDGF-AB and PDGF-BB at 30 ng/ml on mesangial cells.

Table 5

CPM [³H] Thymidine
Incorporated

<u>Sample</u>	<u>Mean</u>	<u>St. dev.</u>
PDGF-AA, 30 ng/ml	492	120
PDGF-AB, 30 ng/ml	1554	114
PDGF-BB, 30 ng/ml	1852	464
Zveg3 GFD, 3 ng/ml	1321	91
Zveg3 GFD, 10 ng/ml	1615	325
Zveg3 GFD, 30 ng/ml	1545	237
Zveg3 GFD, 100 ng/ml	1677	88
10% FBS	1447	174
Buffer control	392	109

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration,

various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

1. A method of determining a value of a function of a variable, the method comprising: receiving a value of the variable; and determining the value of the function of the variable based on the received value of the variable.

CLAIMS

What is claimed is:

1. A method of reducing cell proliferation or extracellular matrix production in a mammal comprising administering to the mammal a composition comprising a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides, in an amount sufficient to reduce cell proliferation or extracellular matrix production.
2. The method of claim 1 wherein proliferation of mesangial, endothelial, smooth muscle, fibroblast, osteoblast, osteoclast, stellate, or interstitial cells is reduced.
3. The method of claim 1 wherein extracellular matrix production is reduced.
4. The method of claim 1 wherein the mammal is suffering from a fibroproliferative disorder of the liver.
5. The method of claim 1 wherein the mammal is suffering from a fibroproliferative disorder of the kidney.
6. The method of claim 1 wherein the mammal is suffering from a fibroproliferative disorder of bone.
7. The method of claim 1 wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies and inhibitory polynucleotides.
8. The method of claim 7 wherein the antagonist is an anti-zveg3 antibody.
9. The method of claim 8 wherein the antibody is a monoclonal antibody.
10. The method of claim 7 wherein the antagonist is selected from the group consisting of antisense polynucleotides, ribozyme-encoding polynucleotides, and external guide sequence-encoding polynucleotides.

11. A method of treating fibrosis in a mammal comprising administering to the mammal a composition comprising a therapeutically effective amount of a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides.

12. The method of claim 11 wherein the fibrosis is liver fibrosis.

13. The method of claim 11 wherein the fibrosis is kidney fibrosis.

14. The method of claim 11 wherein the antagonist is an anti-zveg3 antibody.

15. The method of claim 14 wherein the antibody is a monoclonal antibody.

16. The method of claim 11 wherein the antagonist is selected from the group consisting of antisense polynucleotides, ribozyme-encoding polynucleotides, and external guide sequence-encoding polynucleotides.

17. A method of reducing stellate cell activation in a mammal comprising administering to the mammal a composition comprising a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle, wherein the zveg3 antagonist is selected from the group consisting of anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides, in an amount sufficient to reduce stellate cell activation.


18. The method of claim 17 wherein the stellate cells are liver stellate cells.

METHOD OF TREATING FIBROSIS

ABSTRACT OF THE DISCLOSURE

Materials and methods for treating fibrosis in a mammal are disclosed. The methods comprise administering to a mammal a composition comprising a therapeutically effective amount of a zveg3 antagonist in combination with a pharmaceutically acceptable delivery vehicle. Zveg3 antagonists include anti-zveg3 antibodies, mitogenically inactive receptor-binding zveg3 variant polypeptides, and inhibitory polynucleotides. Within one embodiment of the invention the fibrosis is liver fibrosis.

00941030



50	-1.10	=====	I
51	-0.80	=====	t
52	-0.18	==	V
53	-0.23	==	S
54	-0.37	=====	T
55	-0.25	=====	N
56	-0.28	=====	G
57	0.22		S ==
58	-0.25	=====	I
59	0.05		H
60	0.05		S
61	-0.07	=	P
62	-0.05		R
63	-0.55	=====	F
64	0.37		P =====
65	0.40		H =====
66	0.42		T =====
67	0.23		Y ==
68	-0.08	=	P
69	-0.33	=====	R
70	-1.40	=====	N
71	-0.93	=====	T
72	-1.17	=====	V
73	-1.17	=====	L
74	-0.95	=====	V
75	-0.95	=====	W
76	0.12		R =
77	0.12		L =
78	0.45		V =====
79	0.45		A =====
80	-0.03		V
81	-0.08	=	E
82	-0.55	=====	E
83	-1.35	=====	N
84	-1.45	=====	V
85	-1.62	=====	W
86	-0.55	=====	I
87	0.25		Q ===
88	0.72		L =====
89	0.60		T =====
90	0.67		F =====
91	0.78		D =====
92	0.78		E =====
93	0.78		R =====
94	0.28		F ==
95	1.20		G =====
96	1.70		L =====
97	2.50		E =====
98	1.70		D =====
99	1.03		P =====
100	1.53		E =====
101	0.65		D =====
102	0.65		D =====

Fig. 1B

103	-0.27	====	I
104	-0.22	==	C
105	0.45		K =====
106	-0.30	====	Y
107	0.58		D =====
108	0.58		F =====
109	1.00		V =====
110	1.30		E =====
111	1.30		V =====
112	1.55		E =====
113	0.98		E =====
114	0.18		P ==
115	-0.12	=	S
116	-0.17	==	D
117	-0.17	==	G
118	-0.73	=====	T
119	-0.83	=====	I
120	-0.53	=====	L
121	-0.80	=====	G
122	-0.80	=====	R
123	-1.37	=====	W
124	-1.05	=====	C
125	-0.88	=====	G
126	-0.88	=====	s
127	0.18		G ==
128	0.22		T ==
129	-0.02		V
130	0.28		P ===
131	0.78		G =====
132	0.78		K =====
133	0.32		Q ===
134	0.32		I ===
135	0.32		S ===
136	0.77		K =====
137	-0.03		G
138	0.47		N =====
139	0.02		Q
140	-0.27	====	I
141	0.08		R =
142	0.08		I =
143	0.88		R =====
144	0.00		F
145	0.00		V
146	0.25		S ===
147	0.25		D ===
148	0.25		E ===
149	-0.25	====	Y
150	0.13		F =
151	0.13		P =
152	-0.03		S
153	-0.38	====	E
154	-0.97	=====	P
155	-1.35	=====	G
156	-1.32	=====	F

Fig. 1C

157	-1.20	=====	C
158	-1.28	=====	I
159	-1.20	=====	H
160	-1.12	=====	Y
161	-0.70	=====	N
162	-1.15	=====	I
163	-0.92	=====	V
164	-0.17	==	M
165	-0.03		P
166	-0.28	===	Q
167	-0.27	===	F
168	0.15		T =
169	0.27		E ===
170	-0.48	=====	A
171	-0.70	=====	V
172	-0.45	=====	S
173	-0.50	=====	P
174	-0.45	=====	S
175	-0.58	=====	V
176	-0.63	=====	L
177	-0.33	===	P
178	-0.63	=====	P
179	-0.13	=	S
180	-0.48	=====	A
181	-0.70	=====	L
182	-0.37	=====	P
183	-0.33	=====	L
184	-0.12	=	D
185	-0.92	=====	L
186	-1.18	=====	L
187	-0.97	=====	N
188	-1.42	=====	N
189	-2.02	=====	A
190	-2.00	=====	I
191	-2.00	=====	t
192	-0.93	=====	A
193	-0.35	=====	F
194	-0.23	==	S
195	0.03		T
196	0.60		L =====
197	0.52		E =====
198	-0.28	===	D
199	-0.28	===	L
200	0.02		I
201	0.82		R =====
202	0.82		Y =====
203	1.03		L =====
204	1.37		E =====
205	0.57		P =====
206	1.07		E =====
207	0.27		R ===
208	0.27		w ===
209	0.93		Q =====
210	0.60		L =====


Fig. 1D

211	0.52	D	=====
212	0.52	L	=====
213	0.82	E	=====
214	0.25	D	=====
215	-0.42	=====	L
216	-0.08	=	Y
217	0.00		R
218	-0.80	=====	P
219	-0.80	=====	T
220	-0.23	==	w
221	-0.15	=	Q
222	-0.60	=====	L
223	-0.55	=====	L
224	-0.67	=====	G
225	-0.67	=====	K
226	-0.67	=====	A
227	-0.08	=	F
228	0.38	V	=====
229	1.13	F	=====
230	1.30	G	=====
231	1.05	R	=====
232	1.05	K	=====
233	0.25	S	=====
234	0.23	R	=====
235	-0.57	=====	V
236	-0.62	=====	V
237	-0.43	=====	D
238	-0.43	=====	L
239	0.37	N	=====
240	0.08	L	=====
241	0.88	L	=====
242	0.88	T	=====
243	0.57	E	=====
244	0.12	E	=====
245	-0.55	=====	V
246	-0.37	=====	R
247	-0.87	=====	L
248	-0.07	=	Y
249	0.35	S	=====
250	-0.12	=	C
251	0.10	T	=====
252	-0.08	=	P
253	-0.03	R	=====
254	-0.83	=====	N
255	-0.37	=====	F
256	0.55	S	=====
257	1.00	V	=====
258	0.95	S	=====
259	1.40	I	=====
260	2.20	R	=====
261	1.63	E	=====
262	1.63	E	=====
263	1.07	L	=====
264	1.07	K	=====

Fig. 1E

265	0.15		R =
266	-0.92	=====	T
267	-0.85	=====	D
268	-1.35	=====	T
269	-1.45	=====	I
270	-1.45	=====	F
271	-1.33	=====	W
272	-1.02	=====	P
273	-0.52	=====	G
274	-0.02		C
275	-0.02		L
276	0.28		L ==
277	0.58		V =====
278	0.87		K =====
279	0.20		R ==
280	-0.38	=====	C
281	-0.38	=====	G
282	-0.55	=====	G
283	-0.85	=====	N
284	-0.97	=====	C
285	-0.77	=====	A
286	-0.85	=====	C
287	-0.65	=====	C
288	0.02		L
289	0.15		H ==
290	0.27		N ==
291	0.07		C =
292	-0.02		N
293	-0.05		E
294	-0.50	=====	C
295	0.17		Q ==
296	-0.12	=	C
297	-0.02		V
298	0.73		P =====
299	1.23		S =====
300	1.20		K =====
301	0.62		V =====
302	1.37		T =====
303	1.18		K =====
304	0.38		K =====
305	-0.08	=	Y
306	-0.40	=====	H
307	0.18		E ==
308	-0.32	=====	V
309	0.43		L =====
310	0.67		Q =====
311	0.63		L =====
312	0.68		R =====
313	0.68		P =====
314	0.68		K =====
315	-0.12	=	T
316	-0.13	=	G
317	0.37		V =====
318	0.67		R =====

Fig. 1F

[illegible]

MSLFGLLLLTALAGQRQGTQAESNLSSKFQFSSNKEQNGVQDPQHERIITVSTNGSIHS
 MLLLGLLLLLTALAGQRTGTRAESNLSSKLQLSSDKEQNGVQDPRHERVVTISGNGSIHS
 10 20 30 40 50 60

PRFPHTYPRNTVLVWRLVAVEENVWIQLTFDERFGLEDPEDDICKYDFVEVEEPSDGTIL
 PKFPHTYPRNMVLVWRLVAVDENVRIQLTFDERFGLEDPEDDICKYDFVEVEEPSDGSVL
 70 80 90 100 110 120

GRWCGSGTVPGKQISKGNQIRIRFVSDEYFPSEPGFCIHYNIVMPQFTEAVSPSVLPSSA
 GRWCGSGTVPGKQTSKGNHIRIRFVSDEYFPSEPGFCIHYSIIMPQVTETTSPSVLPSS
 130 140 150 160 170 180

LPLDLLNNAITAFSTLEDLIRYLEPERWQLDLEDLYRPTWQLLGKAFVFGRKSRVVDLNL
 LSLDLLNNAVTAFSTLEELIRYLEPDRWQVDLDSLYKPTWQLLGKAFLYGKKSKVVNLNL
 190 200 210 220 230 240

LTEEVRLYSCTPRNFSVSIREELKRTDTIFWPGCLLVKRCGGNCACCLHNCNECQCVPSK
 LKEEVKLYSCTPRNFSVSIREELKRTDTIFWPGCLLVKRCGGNCACCLHNCNECQCVPRK
 250 260 270 280 290 300

VTKKYHEVLQLRPKTGVRGLHKSLTDVALEHHEECDVCVRGSTGG (SEQ ID NO:2)
 VTKKYHEVLQLRPKTGVKGLHKSLTDVALEHHEECDVCVRGNAGG (SEQ ID NO:4)
 310 320 330 340

Fig. 2

COMBINED DECLARATION FOR PATENT AND POWER OF ATTORNEY (Includes Reference to PCT International Applications)			File No. 00-53	
<p>As a below named inventor, I hereby declare that: My residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:</p> <p>METHOD FOR TREATING FIBROSIS</p> <p>the specification of which (check only one item below):</p> <p><input checked="" type="checkbox"/> is attached hereto <input type="checkbox"/> was filed as United States application Serial No. on October 23, 2000</p> <p>and was amended on _____</p> <p><input type="checkbox"/> was filed as PCT international application Number _____ on _____</p> <p>I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, 1.56. I hereby claim foreign priority benefits under Title 35, United States Code, 119 of any foreign application(s) for patent or inventor's certificate(s) or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate(s) or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:</p>				
PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:				
COUNTRY	APPLICATION NUMBER	DATE OF FILING	PRIORITY CLAIMED	
			<input type="checkbox"/> YES <input type="checkbox"/> NO	
			<input type="checkbox"/> YES <input type="checkbox"/> NO	
			<input type="checkbox"/> YES <input type="checkbox"/> NO	
I hereby claim the benefit under Title 35 United States Code 119(e) of any United States provisional application(s) listed below.				
U.S. APPLICATION NUMBER		U.S. FILING DATE		
60/161,653		October 21, 1999		
60/165,255		November 12, 1999		
60/222,223		August 1, 2000		
I hereby claim the benefit under Title 35, United States Code 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, 1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:				
PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT				
U.S. APPLICATIONS		STATUS (check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	Patented	Pending	Abandoned
PCT APPLICATIONS DESIGNATING THE U.S.				
APPLICATION	FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)		

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Robyn Adams
Reg. No. 44,495

Jennifer K. Johnson
Reg. No. 43,696

Phillip B.C. Jones
Reg. No. 38,195

Debra K. Leith
Reg. No. 32,619

Susan E. Lingenfelter
Reg. No. 41,156

Paul G. Lunn
Reg. No. 32,743

Gary E. Parker
Reg. No. 31,648

Deborah A. Sawislak
Reg. No. 37,438

Send Correspondence To: Gary E. Parker
ZymoGenetics, Inc.
1201 Eastlake Avenue East
Seattle, WA 98102

Direct Telephone Calls To:
Gary E. Parker
(206) 442-6673

1	Full Name	Family Name Gilbertson	First Given Name Debra	Second Given Name G.
	Residence	City Seattle	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 4025 NE 195th PL	City Seattle	State & Zip Code/Country WA 98155/US
2	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
3	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
4	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
5	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
6	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country

I hereby declare that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3
Date	Date	Date
Signature of Inventor 4	Signature of Inventor 5	Signature of Inventor 6
Date	Date	Date

SEQUENCE LISTING

<110> Gilbertson, Debra G.

<120> METHOD OF TREATING FIBROSIS

<130> 00-53

<160> 18

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1760

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (154)...(1191)

<400> 1

attatgtgga aactaccctg cgattctctg ctgccagagc aggctcggcg cttccacccc	60
agtgcagcct tcccctggcg gtggtgaaag agactcggga gtcgctgctt ccaaagtgcc	120
cgccgtgagt gagctctcac cccagtcagc caa atg agc ctc ttc ggg ctt ctc	174
Met Ser Leu Phe Gly Leu Leu	
1 5	
ctg ctg aca tct gcc ctg gcc ggc cag aga cag ggg act cag gcg gaa	222
Leu Leu Thr Ser Ala Leu Ala Gly Gln Arg Gln Gly Thr Gln Ala Glu	
10 15 20	
tcc aac ctg agt agt aaa ttc cag ttt tcc agc aac aag gaa cag aac	270
Ser Asn Leu Ser Ser Lys Phe Gln Phe Ser Ser Asn Lys Glu Gln Asn	
25 30 35	
gga gta caa gat cct cag cat gag aga att att act gtg tct act aat	318
Gly Val Gln Asp Pro Gln His Glu Arg Ile Ile Thr Val Ser Thr Asn	
40 45 50 55	
gga agt att cac agc cca agg ttt cct cat act tat cca aga aat acg	366
Gly Ser Ile His Ser Pro Arg Phe Pro His Thr Tyr Pro Arg Asn Thr	
60 65 70	

gtc	ttg	gta	tgg	aga	tta	gta	gca	gta	gag	gaa	aat	gta	tgg	ata	caa	414
Val	Leu	Val	Trp	Arg	Leu	Val	Ala	Val	Glu	Glu	Asn	Val	Trp	Ile	Gln	
75				80				85								
ctt	acg	ttt	gat	gaa	aga	ttt	ggg	ctt	gaa	gac	cca	gaa	gat	gac	ata	462
Leu	Thr	Phe	Asp	Glu	Arg	Phe	Gly	Leu	Glu	Asp	Pro	Glu	Asp	Asp	Ile	
90				95				100								
tgc	aag	tat	gat	ttt	gta	gaa	gtt	gag	gaa	ccc	agt	gat	gga	act	ata	510
Cys	Lys	Tyr	Asp	Phe	Val	Glu	Val	Glu	Glu	Pro	Ser	Asp	Gly	Thr	Ile	
105				110				115								
tta	ggg	cgc	tgg	tgt	ggg	tct	ggg	act	gta	cca	gga	aaa	cag	att	tct	558
Leu	Gly	Arg	Trp	Cys	Gly	Ser	Gly	Thr	Val	Pro	Gly	Lys	Gln	Ile	Ser	
120				125				130				135				
aaa	gga	aat	caa	att	agg	ata	aga	ttt	gta	tct	gat	gaa	tat	ttt	cct	606
Lys	Gly	Asn	Gln	Ile	Arg	Ile	Arg	Phe	Val	Ser	Asp	Glu	Tyr	Phe	Pro	
140				145				150								
tct	gaa	cca	ggg	ttc	tgc	atc	cac	tac	aac	att	gtc	atg	cca	caa	ttc	654
Ser	Glu	Pro	Gly	Phe	Cys	Ile	His	Tyr	Asn	Ile	Val	Met	Pro	Gln	Phe	
155				160				165								
aca	gaa	gct	gtg	agt	cct	tca	gtg	cta	ccc	cct	tca	gct	ttg	cca	ctg	702
Thr	Glu	Ala	Val	Ser	Pro	Ser	Val	Leu	Pro	Pro	Ser	Ala	Leu	Pro	Leu	
170				175				180								
gac	ctg	ctt	aat	aat	gct	ata	act	gcc	ttt	agt	acc	ttg	gaa	gac	ctt	750
Asp	Leu	Leu	Asn	Asn	Ala	Ile	Thr	Ala	Phe	Ser	Thr	Leu	Glu	Asp	Leu	
185				190				195								
att	cga	tat	ctt	gaa	cca	gag	aga	tgg	cag	ttg	gac	tta	gaa	gat	cta	798
Ile	Arg	Tyr	Leu	Glu	Pro	Glu	Arg	Trp	Gln	Leu	Asp	Leu	Glu	Asp	Leu	
200				205				210				215				
tat	agg	cca	act	tgg	caa	ctt	ctt	ggc	aag	gct	ttt	gtt	ttt	gga	aga	846
Tyr	Arg	Pro	Thr	Trp	Gln	Leu	Leu	Gly	Lys	Ala	Phe	Val	Phe	Gly	Arg	
220				225				230								
aaa	tcc	aga	gtg	gtg	gat	ctg	aac	ctt	cta	aca	gag	gag	gta	aga	tta	894
Lys	Ser	Arg	Val	Val	Asp	Leu	Asn	Leu	Leu	Thr	Glu	Glu	Val	Arg	Leu	
235				240				245								

```
<210> 2
<211> 345
<212> PRT
<213> Homo sapiens
```

 $\langle 400 \rangle$ 2

Met Ser Leu Phe Gly Leu Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln
 1 5 10 15
 Arg Gln Gly Thr Gln Ala Glu Ser Asn Leu Ser Ser Lys Phe Gln Phe
 20 25 30
 Ser Ser Asn Lys Glu Gln Asn Gly Val Gln Asp Pro Gln His Glu Arg
 35 40 45
 Ile Ile Thr Val Ser Thr Asn Gly Ser Ile His Ser Pro Arg Phe Pro
 50 55 60
 His Thr Tyr Pro Arg Asn Thr Val Leu Val Trp Arg Leu Val Ala Val
 65 70 75 80
 Glu Glu Asn Val Trp Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu
 85 90 95
 Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu
 100 105 110
 Glu Pro Ser Asp Gly Thr Ile Leu Gly Arg Trp Cys Gly Ser Gly Thr
 115 120 125
 Val Pro Gly Lys Gln Ile Ser Lys Gly Asn Gln Ile Arg Ile Arg Phe
 130 135 140
 Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr
 145 150 155 160
 Asn Ile Val Met Pro Gln Phe Thr Glu Ala Val Ser Pro Ser Val Leu
 165 170 175
 Pro Pro Ser Ala Leu Pro Leu Asp Leu Leu Asn Asn Ala Ile Thr Ala
 180 185 190
 Phe Ser Thr Leu Glu Asp Leu Ile Arg Tyr Leu Glu Pro Glu Arg Trp
 195 200 205
 Gln Leu Asp Leu Glu Asp Leu Tyr Arg Pro Thr Trp Gln Leu Leu Gly
 210 215 220
 Lys Ala Phe Val Phe Gly Arg Lys Ser Arg Val Val Asp Leu Asn Leu
 225 230 235 240
 Leu Thr Glu Glu Val Arg Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser
 245 250 255
 Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro
 260 265 270
 Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu
 275 280 285
 His Asn Cys Asn Glu Cys Gln Cys Val Pro Ser Lys Val Thr Lys Lys
 290 295 300
 Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Arg Gly Leu
 305 310 315 320
 His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp
 325 330 335
 Cys Val Cys Arg Gly Ser Thr Gly Gly
 340 345

<210> 3
 <211> 3571
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1049)...(2086)

<400> 3

gaattcccg gtcgacccac gcgtccgggc gccagggga aaggaagctg ggggccgcct 60
 ggcggcattc ctcgccgag tgtgggctcc gtctgccgcg gggcccgag tgccccctgt 120
 ctgcgccagc acctgttggc ccgccagctg gccgccgcg cccccgcgc cccccgcgcc 180
 cgccggccg ccagccccgc gcccgcgcg ccgccgctg ggggaaagt gagacgggga 240
 ggggacaaga gcgacccctc aggccagcca ggccttccct tagccgcccg tgcttagccg 300
 ccacctctcc tcagccctgc gtctgccct gccttagggc aggcaccca gcgctcgcga 360
 ctccgagccg cccaagctct ccgcgttcc cgcagcactt cgccggtacc cgagggaact 420
 tcggtggcca ccgactgcag caaggaggag gctccgcggt ggatccgggc cagtcccag 480
 tcgtccccgc ggcctctctg ccgcccggg accgcgcgcg cactcgcagg gcacggtccc 540
 ctccccccag gtgggggtgg ggcgcgcct gccgccccga tcagcagctt tgtcattgat 600
 cccaaggtgc tcgctctgct gccgacctg cttccagtct ggcttggcg gaccccgagt 660
 cctgcctgt gtctgtccc ccaaactgac aggtgctccc tgcgagtcgc cagactcat 720
 cgccgctccc ccggtcccc accccttctt tcctccctcg cctaccccca cccccgcac 780
 ttccgacag ctcaggattt gtttaaact tgggaaactg gttcagggtc aggttttgct 840
 ttgatccttt tcaaaaactg gagacacaga agagggtctt aggaaaaact tttgatggg 900
 attatgtgga aactaccctg cgattctctg ctgccagagc cggccaggcg cttccaccgc 960
 agcgcagcct ttccccggt gggctgagcc ttggagtcgt cgcttcccca gtgcccgcg 1020
 cgagtgcgac ctcgccccag tcagccaa atg ctc ctc ctc ggc ctc ctc ctg 1072

Met Leu Leu Leu Gly Leu Leu Leu

1

5

ctg aca tct gcc ctg gcc ggc caa aga acg ggg act cgg gct gag tcc 1120
 Leu Thr Ser Ala Leu Ala Gly Gln Arg Thr Gly Thr Arg Ala Glu Ser
 10 15 20

aac ctg agc agc aag ttg cag ctc tcc agc gac aag gaa cag aac gga 1168
 Asn Leu Ser Ser Lys Leu Gln Leu Ser Ser Asp Lys Glu Gln Asn Gly
 25 30 35 40

gtg caa gat ccc cgg cat gag aga gtt gtc act ata tct ggt aat ggg 1216
 Val Gln Asp Pro Arg His Glu Arg Val Val Thr Ile Ser Gly Asn Gly
 45 50 55

agc atc cac agc ccg aag ttt cct cat aca tac cca aga aat atg gtg Ser Ile His Ser Pro Lys Phe Pro His Thr Tyr Pro Arg Asn Met Val	1264
60 65 70	
ctg gtg tgg aga tta gtt gca gta gat gaa aat gtg cgg atc cag ctg Leu Val Trp Arg Leu Val Ala Val Asp Glu Asn Val Arg Ile Gln Leu	1312
75 80 85	
aca ttt gat gag aga ttt ggg ctg gaa gat cca gaa gac gat ata tgc Thr Phe Asp Glu Arg Phe Gly Leu Glu Asp Pro Glu Asp Asp Ile Cys	1360
90 95 100	
aag tat gat ttt gta gaa gtt gag gag ccc agt gat gga agt gtt tta Lys Tyr Asp Phe Val Glu Val Glu Glu Pro Ser Asp Gly Ser Val Leu	1408
105 110 115 120	
gga cgc tgg tgt ggt tct ggg act gtg cca gga aag cag act tct aaa Gly Arg Trp Cys Gly Ser Gly Thr Val Pro Gly Lys Gln Thr Ser Lys	1456
125 130 135	
gga aat cat atc agg ata aga ttt gta tct gat gag tat ttt cca tct Gly Asn His Ile Arg Ile Arg Phe Val Ser Asp Glu Tyr Phe Pro Ser	1504
140 145 150	
gaa ccc gga ttc tgc atc cac tac agt att atc atg cca caa gtc aca Glu Pro Gly Phe Cys Ile His Tyr Ser Ile Ile Met Pro Gln Val Thr	1552
155 160 165	
gaa acc acg agt cct tcg gtg ttg ccc cct tca tct ttg tca ttg gac Glu Thr Thr Ser Pro Ser Val Leu Pro Pro Ser Ser Leu Ser Leu Asp	1600
170 175 180	
ctg ctc aac aat gct gtg act gcc ttc agt acc ttg gaa gag ctg att Leu Leu Asn Asn Ala Val Thr Ala Phe Ser Thr Leu Glu Glu Leu Ile	1648
185 190 195 200	
cgg tac cta gag cca gat cga tgg cag gtg gac ttg gac agc ctc tac Arg Tyr Leu Glu Pro Asp Arg Trp Gln Val Asp Leu Asp Ser Leu Tyr	1696
205 210 215	
aag cca aca tgg cag ctt ttg ggc aag gct ttc ctg tat ggg aaa aaa Lys Pro Thr Trp Gln Leu Leu Gly Lys Ala Phe Leu Tyr Gly Lys Lys	1744
220 225 230	

agc Ser	aaa Lys	gtg Val	gtg Val	aat Asn	ctg Leu	aat Asn	ctc Leu	ctc Leu	aag Lys	gaa Glu	gag Glu	gta Val	aaa Lys	ctc Leu	tac Tyr	1792
235			240			245										
agc Ser	tgc Cys	aca Thr	ccc Pro	cgg Arg	aac Asn	ttc Phe	tca Ser	gtg Val	tcc Ser	ata Ile	cgg Arg	gaa Glu	gag Glu	cta Leu	aag Lys	1840
250			255			260										
agg Arg	aca Thr	gat Asp	acc Thr	ata Ile	ttc Phe	tgg Trp	cca Pro	ggt Gly	tgt Cys	ctc Leu	ctg Leu	gtc Val	aag Lys	cgc Arg	tgt Cys	1888
265			270			275			280							
gga Gly	gga Gly	aat Asn	tgt Cys	gcc Ala	tgt Cys	tgt Cys	ctc Leu	cat His	aat Asn	tgc Cys	aat Asn	gaa Glu	tgt Cys	cag Gln	tgt Cys	1936
285			290			295										
gtc Val	cca Pro	cgt Arg	aaa Lys	gtt Val	aca Thr	aaa Lys	aag Lys	tac Tyr	cat His	gag Glu	gtc Val	ctt Leu	cag Gln	ttg Leu	aga Arg	1984
300			305			310										
cca Pro	aaa Lys	act Thr	gga Gly	gtc Val	aag Lys	gga Gly	ttg Leu	cat His	aag Lys	tca Ser	ctc Leu	act Thr	gat Asp	gtg Val	gct Ala	2032
315			320			325										
ctg Leu	gaa Glu	cac His	cac His	gag Glu	gaa Glu	tgt Cys	gac Asp	tgt Cys	gtg Val	tgt Cys	aga Arg	gga Gly	aac Asn	gca Ala	gga Gly	2080
330			335			340										
ggg Gly	taa *	ctgcagcctt	cgtagcagca	cacgtgagca	ctggcattct	gtgtaccccc										2136
345																
acaagcaacc	ttcatcccca	ccagcgttgg	ccgcagggct	ctcagctgct	gatgctggct											2196
atggtaaaga	tcttactcgt	ctccaaccaa	attctcagtt	gtttgcttca	atagccttcc											2256
cctgcaggac	ttcaagtgtc	ttctaaaaga	ccagaggcac	caagaggagt	caatcacaaa											2316
gcactgcctt	ctagaggaag	cccagacaat	ggtcttctga	ccacagaaac	aatgaaatg											2376
aatgtagatc	gctagcaaac	tctggagtga	cagcatttct	tttccactga	cagaatgggtg											2436
tagcttagtt	gtcttgatat	gggcaagtga	tgtcagcaca	agaaaatgggt	gaaaaacaca											2496
cacttgattg	tgaacaatgc	agaaataactt	ggatttctcc	aacctgtttg	catagataga											2556
cagatgctct	gttttctaca	aactcaaagc	ttttagagag	cagctatggt	aataggaatt											2616
aatgtgcca	tgctgaaagg	aaagactgaa	gttttcaatg	cttggaact	tctccgcaat											2676
ttggaggaaa	ggtgcgggtca	tggtttggag	aaagcacacc	tgcacagagg	agtggccttc											2736
ccttccttc	cctctgaggt	ggcttctgtg	tttcattgtg	tatatatttta	tattctcctt											2796
ttgacattat	aactgttggc	ttttctaata	ttgttaaata	tttctatttt	taccaaaggt											2856

```

atttaatat cttttttatg acaacctaga gcaattat ttagcttgat aatTTTTTTT 2916
tctaaacaaa attgttatag ccagaagaac aaagatgatt gatataaaaa tcttggtgct 2976
ctgacaaaaa catatgtatt tcttccttgt atgggtgctag agcttagcgt catctgcatt 3036
tgaaaagatg gaatggggaa gtttttagaa ttggtaggtc gcagggacag tttgataaca 3096
actgtactat catcaattcc caattctgtt cttagagcta cgaacagaac agagcttgag 3156
taaatatgga gccattgcta acctaccctt ttctatggga aataggagta tagctcagag 3216
aagcacgtcc ccagaaacct cgaccatttc taggcacagt gttctgggct atgctgcgct 3276
gtatggacat atcctattta tttcaatact agggttttat tacctttaaa ctctgctcca 3336
tacacttgta ttaatacatg gatattttta tgtacagaag tatatcattt aaggagttca 3396
cttattatac tctttggcaa ttgcaaagaa aatcaacata atacattgct tgtaaagtct 3456
taatctgtgc ccaagttttg tgggtgactat ttgaattaaa atgtattgaa tcatcaaata 3516
aaataatctg gctattttgg ggaaaaaaaa aaaaaaaaaa aaaaagggcg gccgc 3571

```

<210> 4

<211> 345

<212> PRT

<213> Mus musculus

<400> 4

```

Met Leu Leu Leu Gly Leu Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln
 1          5          10          15
Arg Thr Gly Thr Arg Ala Glu Ser Asn Leu Ser Ser Lys Leu Gln Leu
      20          25          30
Ser Ser Asp Lys Glu Gln Asn Gly Val Gln Asp Pro Arg His Glu Arg
      35          40          45
Val Val Thr Ile Ser Gly Asn Gly Ser Ile His Ser Pro Lys Phe Pro
      50          55          60
His Thr Tyr Pro Arg Asn Met Val Leu Val Trp Arg Leu Val Ala Val
      65          70          75          80
Asp Glu Asn Val Arg Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu
      85          90          95
Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu
      100         105         110
Glu Pro Ser Asp Gly Ser Val Leu Gly Arg Trp Cys Gly Ser Gly Thr
      115         120         125
Val Pro Gly Lys Gln Thr Ser Lys Gly Asn His Ile Arg Ile Arg Phe
      130         135         140
Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr
      145         150         155         160
Ser Ile Ile Met Pro Gln Val Thr Glu Thr Thr Ser Pro Ser Val Leu
      165         170         175
Pro Pro Ser Ser Leu Ser Leu Asp Leu Leu Asn Asn Ala Val Thr Ala
      180         185         190

```

```
<210> 5
<211> 370
<212> PRT
<213> Homo sapiens
```

<400> 5															
Met	His	Arg	Leu	Ile	Phe	Val	Tyr	Thr	Leu	Ile	Cys	Ala	Asn	Phe	Cys
1				5					10					15	
Ser	Cys	Arg	Asp	Thr	Ser	Ala	Thr	Pro	Gln	Ser	Ala	Ser	Ile	Lys	Ala
			20					25					30		
Leu	Arg	Asn	Ala	Asn	Leu	Arg	Arg	Asp	Glu	Ser	Asn	His	Leu	Thr	Asp
		35					40					45			
Leu	Tyr	Arg	Arg	Asp	Glu	Thr	Ile	Gln	Val	Lys	Gly	Asn	Gly	Tyr	Val
	50					55					60				
Gln	Ser	Pro	Arg	Phe	Pro	Asn	Ser	Tyr	Pro	Arg	Asn	Leu	Leu	Leu	Thr
65					70					75					80
Trp	Arg	Leu	His	Ser	Gln	Glu	Asn	Thr	Arg	Ile	Gln	Leu	Val	Phe	Asp
			85						90					95	
Asn	Gln	Phe	Gly	Leu	Glu	Glu	Ala	Glu	Asn	Asp	Ile	Cys	Arg	Tyr	Asp
			100					105					110		
Phe	Val	Glu	Val	Glu	Asp	Ile	Ser	Glu	Thr	Ser	Thr	Ile	Ile	Arg	Gly
		115					120						125		

Gln Leu Asp His His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro
355 360 365
Pro Arg
370

```
<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide primer

<400> 6
```

20

<213> Artificial Sequence

<223> oligonucleotide primer

21

<213> Artificial Sequence

<223> oligonucleotide primer

21

<213> Artificial Sequence

<223> oligonucleotide primer

20

<213> Artificial Sequence

<223> oligonucleotide primer

<400> 10

25

<211> 25

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> oligonucleotide primer

<400> 11

aaactaccct gcgattctct gctgc

25

<210> 12

<211> 21

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> oligonucleotide primer

<400> 12

ggtaaattgga gcttggctga g

21

<210> 13

<211> 25

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> oligonucleotide primer

<400> 13

tctggacgtc ctctgctgg tatag

25

<210> 14

<211> 25

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> oligonucleotide primer

<400> 14

25

<211> 27

<213> Artificial Sequence

<223> oligonucleotide primer

gagtggcaac ttccagggcc aggagag

27

<211> 27

<213> Artificial Sequence

<223> oligonucleotide primer

cttttgctag cctcaaccct gactatc

27

<211> 35

<213> Artificial Sequence

<223> oligonucleotide primer ZC20,180

cgcgcggttt aaacgccacc atgagcctct tcggg

35

<211> 32

<213> Artificial Sequence

<223> oligonucleotide primer ZC20,181

<400> 18

cgtatcggcg cgcctatcc tcctgtgctc cc

008204" P3T36960